

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/55, 9/20, 1/21, 5/10		A2	(11) International Publication Number: WO 99/55883
			(43) International Publication Date: 4 November 1999 (04.11.99)
(21) International Application Number: PCT/US99/09280			(81) Designated States: AE, AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 29 April 1999 (29.04.99)			
(30) Priority Data: 60/083,688 30 April 1998 (30.04.98) US			
(71) Applicant (for all designated States except US): E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): CAHOON, Edgar, B. [US/US]; 2331 West 18th Street, Wilmington, DE 19806 (US). CAHOON, Rebecca, E. [US/US]; 2331 West 18th Street, Wilmington, DE 19806 (US). KINNEY, Anthony, J. [GB/US]; 609 Lore Avenue, Wilmington, DE 19809 (US). RAFALSKI, J., Antoni [US/US]; 2028 Longcome Drive, Wilmington, DE 19810 (US).			
(74) Agent: MAJARIAN, William, R.; E.I. du Pont de Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).			Published Without international search report and to be republished upon receipt of that report.

(54) Title: TRIACYLGLYCEROL LIPASES

```
SEQ ID NO: 37  * * * * *
SEQ ID NO: 38  * * * * *
SEQ ID NO: 39  * * * * *
SEQ ID NO: 40  * * * * *
SEQ ID NO: 41  * * * * *
```

```
SEQ ID NO: 37  * * * * *
SEQ ID NO: 38  * * * * *
SEQ ID NO: 39  * * * * *
SEQ ID NO: 40  * * * * *
SEQ ID NO: 41  * * * * *
```

```
SEQ ID NO: 37  * * * * *
SEQ ID NO: 38  * * * * *
SEQ ID NO: 39  * * * * *
SEQ ID NO: 40  * * * * *
SEQ ID NO: 41  * * * * *
```

```
SEQ ID NO: 37  * * * * *
SEQ ID NO: 38  * * * * *
SEQ ID NO: 39  * * * * *
SEQ ID NO: 40  * * * * *
SEQ ID NO: 41  * * * * *
```

```
SEQ ID NO: 37  * * * * *
SEQ ID NO: 38  * * * * *
SEQ ID NO: 39  * * * * *
SEQ ID NO: 40  * * * * *
SEQ ID NO: 41  * * * * *
```

```
SEQ ID NO: 37  * * * * *
SEQ ID NO: 38  * * * * *
SEQ ID NO: 39  * * * * *
SEQ ID NO: 40  * * * * *
SEQ ID NO: 41  * * * * *
```

```
SEQ ID NO: 37  * * * * *
SEQ ID NO: 38  * * * * *
SEQ ID NO: 39  * * * * *
SEQ ID NO: 40  * * * * *
SEQ ID NO: 41  * * * * *
```

```
SEQ ID NO: 37  * * * * *
SEQ ID NO: 38  * * * * *
SEQ ID NO: 39  * * * * *
SEQ ID NO: 40  * * * * *
SEQ ID NO: 41  * * * * *
```

```
SEQ ID NO: 37  * * * * *
SEQ ID NO: 38  * * * * *
SEQ ID NO: 39  * * * * *
SEQ ID NO: 40  * * * * *
SEQ ID NO: 41  * * * * *
```

```
SEQ ID NO: 37  * * * * *
SEQ ID NO: 38  * * * * *
SEQ ID NO: 39  * * * * *
SEQ ID NO: 40  * * * * *
SEQ ID NO: 41  * * * * *
```

```
SEQ ID NO: 37  * * * * *
SEQ ID NO: 38  * * * * *
SEQ ID NO: 39  * * * * *
SEQ ID NO: 40  * * * * *
SEQ ID NO: 41  * * * * *
```

```
SEQ ID NO: 37  * * * * *
SEQ ID NO: 38  * * * * *
SEQ ID NO: 39  * * * * *
SEQ ID NO: 40  * * * * *
SEQ ID NO: 41  * * * * *
```

(57) Abstract

This invention relates to an isolated nucleic acid fragment encoding a triacylglycerol lipase. The invention also relates to the construction of a chimeric gene encoding all or a portion of the triacylglycerol lipase, in sense or antisense orientation, wherein expression of the chimeric gene results in production of altered levels of the triacylglycerol lipase in a transformed host cell.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE

TRIACYLGLYCEROL LIPASES

This application claims the benefit of U.S. Provisional Application No. 60/083,688, filed April 30, 1998.

FIELD OF THE INVENTION

This invention is in the field of plant molecular biology. More specifically, this invention pertains to nucleic acid fragments encoding triacylglycerol lipases in plants and seeds.

BACKGROUND OF THE INVENTION

True lipases attach triacylglycerols and act at an oil-water interface; they constitute a ubiquitous group of enzymes catalyzing a wide variety of reactions, many with industrial potential. Triacylglycerol lipases catalyze the transformation of triacylglycerol and water into diacylglycerol and a fatty acid anion. Human gastric lipase, rat lingual lipase, and human hepatic lysosomal lipase amino acid sequences are homologous but are unrelated to porcine pancreatic lipase apart from a 6 amino-acid sequence around the essential Ser-152 of porcine pancreatic lipase (Bodmer, M. W. (1987) *Biochim Biophys Acta* 909:237-244). These enzymes are glycosylated, contain a hydrophobic signal peptide, and belong to a gene family of acid lipases (Ameis, D. et al. (1994) *Eur J Biochem* 219:905-914). Lysosomal acid lipase (LAL) is a hydrolase essential for the intracellular degradation of cholesteryl esters and triacylglycerols and participates in the mobilization of seed oil during germination. No plant triacylglycerol lipase cDNAs of this class are currently listed in GenBank.

Neutral triacylglycerol lipases have been widely studied in fungi, bacteria, mammals, and insects. Nucleotide sequences with similarities to neutral triacylglycerol lipases in *Arabidopsis thaliana* and *Ipomea nil* have been described but their function has not yet been proven. The X-ray structure of the *Mucor miehei* triglyceride lipase has been reported, revealing a Ser...His...Asp trypsin-like catalytic triad with an active serine buried under the short helical fragment of a long surface loop (Brady, L. et al. (1990) *Nature* 343:767-770).

It may be useful to isolate triacylglycerol lipase cDNAs from plants that accumulate large amounts of fatty acids with unusual structures. Lacking this ability could be a possible limitation in development of transgenic crops with novel seed oils. Triacylglycerol lipases may also be useful in processing of plant seed oils. Lysosomal acid lipase (LAL) may be used to engineer lipid and cholesteryl ester metabolism and/or lysosome function.

SUMMARY OF THE INVENTION

The instant invention relates to isolated nucleic acid fragments encoding triacylglycerol lipases. Specifically, this invention concerns an isolated nucleic acid fragment encoding an acid or a neutral triacylglycerol lipase. In addition, this invention relates to a nucleic acid fragment that is complementary to the nucleic acid fragment encoding an acid or a neutral triacylglycerol lipase.

An additional embodiment of the instant invention pertains to a polypeptide encoding all or a substantial portion of a triacylglycerol lipase selected from the group consisting of acid and neutral triacylglycerol lipases.

5 In another embodiment, the instant invention relates to a chimeric gene encoding an acid or a neutral triacylglycerol lipase, or to a chimeric gene that comprises a nucleic acid fragment that is complementary to a nucleic acid fragment encoding an acid or a neutral triacylglycerol lipase, operably linked to suitable regulatory sequences, wherein expression of the chimeric gene results in production of levels of the encoded protein in a transformed host cell that is altered (i.e., increased or decreased) from the level produced in an
10 untransformed host cell.

In a further embodiment, the instant invention concerns a transformed host cell comprising in its genome a chimeric gene encoding an acid or a neutral triacylglycerol lipase, operably linked to suitable regulatory sequences. Expression of the chimeric gene results in production of altered levels of the encoded protein in the transformed host cell.
15 The transformed host cell can be of eukaryotic or prokaryotic origin, and include cells derived from higher plants and microorganisms. The invention also includes transformed plants that arise from transformed host cells of higher plants, and seeds derived from such transformed plants.

An additional embodiment of the instant invention concerns a method of altering the
20 level of expression of an acid or a neutral triacylglycerol lipase in a transformed host cell comprising: a) transforming a host cell with a chimeric gene comprising a nucleic acid fragment encoding an or a neutral acid triacylglycerol lipase; and b) growing the transformed host cell under conditions that are suitable for expression of the chimeric gene wherein expression of the chimeric gene results in production of altered levels of acid or
25 neutral triacylglycerol lipase in the transformed host cell.

An addition embodiment of the instant invention concerns a method for obtaining a nucleic acid fragment encoding all or a substantial portion of an amino acid sequence encoding an acid or a neutral triacylglycerol lipase.

30 BRIEF DESCRIPTION OF THE
DRAWINGS AND SEQUENCE DESCRIPTIONS

The invention can be more fully understood from the following detailed description and the accompanying drawings and Sequence Listing which form a part of this application.

Figure 1 depicts the amino acid sequence alignment between the acid triacylglycerol lipase from rice clone rlr72.pk0015.b2 (SEQ ID NO:14), soybean contig assembled from
35 clones sdp3c.pk004.n3 and ssl.pk0022.a1 (SEQ ID NO:18), soybean contig assembled from clones sls1c.pk009.o2, srr1c.pk001.m19 and sre.pk0004.d7 (SEQ ID NO:20), *Canis familiaris* (NCBI General Identifier No. 3041702, SEQ ID NO:35) and *Caenorhabditis elegans* (NCBI General Identifier No. 3165581, SEQ ID NO:36). Amino acids which are conserved among all sequences are indicated with an asterisk (*) while amino acids

conserved only among plant sequences are indicated by a plus sign (+). Dashes are used by the program to maximize alignment of the sequences.

The following sequence descriptions and Sequence Listing attached hereto comply with the rules governing nucleotide and/or amino acid sequence disclosures in patent applications as set forth in 37 C.F.R. §1.821-1.825.

SEQ ID NO:1 is the nucleotide sequence comprising the entire cDNA insert in clone cen3n.pk0129.e9 encoding a portion of a corn acid triacylglycerol lipase.

SEQ ID NO:2 is the deduced amino acid sequence of a portion of a corn acid triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:1.

10 SEQ ID NO:3 is the nucleotide sequence comprising the 3' 647 nucleotides from the cDNA insert in clone ncs.pk0013.h1 encoding the C-terminal quarter of a *Catalpa* acid triacylglycerol lipase

SEQ ID NO:4 is the deduced amino acid sequence of the C-terminal quarter of a *Catalpa* acid triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:3.

15 SEQ ID NO:5 is the nucleotide sequence comprising the 5' 705 nucleotides from the cDNA insert in clone ncs.pk0013.h1 encoding the N-terminal third of a *Catalpa* acid triacylglycerol lipase.

SEQ ID NO:6 is the deduced amino acid sequence of the N-terminal third of a *Catalpa* acid triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:5.

20 SEQ ID NO:7 is the nucleotide sequence comprising the contig assembled from a portion of the cDNA insert in clones p0075.cslag33r, p0126.cnlay46r and p0014.ctuty54r encoding a substantial portion of a corn acid triacylglycerol lipase.

SEQ ID NO:8 is the deduced amino acid sequence of a substantial portion of a corn acid triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:7.

25 SEQ ID NO:9 is the nucleotide sequence comprising a portion of the cDNA insert in clone p0102.ceral64r encoding a portion of a corn acid triacylglycerol lipase.

SEQ ID NO:10 is the deduced amino acid sequence of a portion of a corn acid triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:9.

30 SEQ ID NO:11 is the nucleotide sequence comprising a portion of the cDNA insert in clone p0126.cnlcm37r encoding a portion of a corn acid triacylglycerol lipase.

SEQ ID NO:12 is the deduced amino acid sequence of a portion of a corn acid triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:11.

SEQ ID NO:13 is the nucleotide sequence comprising the entire cDNA insert in clone rlr72.pk0015.b2 encoding an entire rice acid triacylglycerol lipase.

35 SEQ ID NO:14 is the deduced amino acid sequence of an entire rice acid triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:13.

SEQ ID NO:15 is the nucleotide sequence comprising a portion of the cDNA insert in clone rsl1n.pk012.h7 encoding a portion of a rice acid triacylglycerol lipase.

SEQ ID NO:16 is the deduced amino acid sequence of a portion of a rice acid triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:15.

SEQ ID NO:17 is the nucleotide sequence comprising the contig assembled from the entire cDNA insert in clone ssl.pk0022.a1 and a portion of the cDNA insert in clone
5 sdp3c.pk004.n3 encoding an entire soybean acid triacylglycerol lipase.

SEQ ID NO:18 is the deduced amino acid sequence of an entire soybean acid triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:17.

SEQ ID NO:19 is the nucleotide sequence comprising the contig assembled from the entire cDNA insert in clone sre.pk0004.d7 and a portion of the cDNA insert in clones
10 sls1c.pk009.o2 and srr1c.pk001.m19 encoding an entire soybean acid triacylglycerol lipase.

SEQ ID NO:20 is the deduced amino acid sequence of an entire soybean acid triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:19.

SEQ ID NO:21 is the nucleotide sequence comprising the entire cDNA insert in clone cr1n.pk0145.c6 encoding half of a corn neutral triacylglycerol lipase.

15 SEQ ID NO:22 is the deduced amino acid sequence of half of a corn neutral triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:21.

SEQ ID NO:23 is the nucleotide sequence comprising the contig assembled from a portion of the cDNA insert in clones p0010.cbpbe40r, p0083.cldcq17r, p0048.cqlac25r, p0118.chsbw59r, cr1.pk0011.c9 and cdolc.pk002.c22 encoding an entire corn neutral
20 triacylglycerol lipase.

SEQ ID NO:24 is the deduced amino acid sequence of an entire corn neutral triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:23.

SEQ ID NO:25 is the nucleotide sequence comprising the contig assembled from the entire cDNA insert in clone cr1n.pk0127.h8 and a portion of the cDNA insert in clones
25 p0037.crwan02r, p0004.cb1fm22r, p0004.cb1ei43r, cco1n.pk068.o9 and p0093.cssao39r encoding most of a corn neutral triacylglycerol lipase.

SEQ ID NO:26 is the deduced amino acid sequence of most of a corn neutral triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:25.

SEQ ID NO:27 is the nucleotide sequence comprising a portion of the cDNA insert in
30 clone rdr1f.pk002.f11 encoding a portion of a rice neutral triacylglycerol lipase.

SEQ ID NO:28 is the deduced amino acid sequence of a portion of a rice neutral triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:27.

SEQ ID NO:29 is the nucleotide sequence comprising the contig assembled from the entire cDNA insert in clone sre.pk0058.b1 and a portion of the cDNA insert in clone
35 sah1c.pk001.k20 encoding a substantial portion of a soybean neutral triacylglycerol lipase.

SEQ ID NO:30 is the deduced amino acid sequence of a substantial portion of a soybean neutral triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:29.

SEQ ID NO:31 is the nucleotide sequence comprising the entire cDNA insert in clone sr1.pk0079.e1 encoding the C-terminal half of a soybean neutral triacylglycerol lipase.

SEQ ID NO:32 is the deduced amino acid sequence of the C-terminal half of a soybean neutral triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:31.

SEQ ID NO:33 is the nucleotide sequence comprising the entire cDNA insert in clone wr1.pk0115.f5 encoding a portion of a wheat neutral triacylglycerol lipase.

SEQ ID NO:34 is the deduced amino acid sequence of a portion of a wheat neutral triacylglycerol lipase derived from the nucleotide sequence of SEQ ID NO:33.

SEQ ID NO:35 is the amino acid sequence of a *Canis familiaris* acid triacylglycerol lipase, NCBI General Identifier No. 3041702.

SEQ ID NO:36 is the amino acid sequence of a *Caenorhabditis elegans* acid triacylglycerol lipase, NCBI General Identifier No. 3165581.

The Sequence Listing contains the one letter code for nucleotide sequence characters and the three letter codes for amino acids as defined in conformity with the IUPAC-IUBMB standards described in *Nucleic Acids Research* 13:3021-3030 (1985) and in the *Biochemical Journal* 219 (No. 2):345-373 (1984) which are herein incorporated by reference. The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

DETAILED DESCRIPTION OF THE INVENTION

In the context of this disclosure, a number of terms shall be utilized. As used herein, an "isolated nucleic acid fragment" is a polymer of RNA or DNA that is single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases. An isolated nucleic acid fragment in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA or synthetic DNA. As used herein, "contig" refers to an assemblage of overlapping nucleic acid sequences to form one contiguous nucleotide sequence. For example, several DNA sequences can be compared and aligned to identify common or overlapping regions. The individual sequences can then be assembled into a single contiguous nucleotide sequence.

As used herein, "substantially similar" refers to nucleic acid fragments wherein changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the protein encoded by the DNA sequence. "Substantially similar" also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by antisense or co-suppression technology. "Substantially similar" also refers to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially affect the functional properties of the resulting transcript vis-à-vis the ability to mediate alteration of gene expression by antisense or co-suppression technology or alteration of the functional

properties of the resulting protein molecule. It is therefore understood that the invention encompasses more than the specific exemplary sequences.

For example, it is well known in the art that antisense suppression and co-suppression of gene expression may be accomplished using nucleic acid fragments representing less than the entire coding region of a gene, and by nucleic acid fragments that do not share 100% sequence identity with the gene to be suppressed. Moreover, alterations in a gene which result in the production of a chemically equivalent amino acid at a given site, but do not effect the functional properties of the encoded protein, are well known in the art. Thus, a codon for the amino acid alanine, a hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue, such as glycine, or a more hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a functionally equivalent product. Nucleotide changes which result in alteration of the N-terminal and C-terminal portions of the protein molecule would also not be expected to alter the activity of the protein. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products. Moreover, substantially similar nucleic acid fragments may also be characterized by their ability to hybridize, under stringent conditions (0.1X SSC, 0.1% SDS, 65°C), with the nucleic acid fragments disclosed herein.

Substantially similar nucleic acid fragments of the instant invention may also be characterized by the percent similarity of the amino acid sequences that they encode to the amino acid sequences disclosed herein, as determined by algorithms commonly employed by those skilled in this art. Preferred are those nucleic acid fragments whose nucleotide sequences encode amino acid sequences that are 80% similar to the amino acid sequences reported herein. More preferred nucleic acid fragments encode amino acid sequences that are 90% similar to the amino acid sequences reported herein. Most preferred are nucleic acid fragments that encode amino acid sequences that are 95% similar to the amino acid sequences reported herein. Sequence alignments and percent similarity calculations were performed using the Megalign program of the LASARGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins, D. G. and Sharp, P. M. (1989) *CABIOS* 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

A "substantial portion" of an amino acid or nucleotide sequence comprises enough of the amino acid sequence of a polypeptide or the nucleotide sequence of a gene to afford putative identification of that polypeptide or gene, either by manual evaluation of the sequence by one skilled in the art, or by computer-automated sequence comparison and

identification using algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al. (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/). In general, a sequence of ten or more contiguous amino acids or thirty or more nucleotides is necessary in order to putatively identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene specific oligonucleotide probes comprising 20-30 contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., *in situ* hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12-15 bases may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a "substantial portion" of a nucleotide sequence comprises enough of the sequence to afford specific identification and/or isolation of a nucleic acid fragment comprising the sequence. The instant specification teaches partial or complete amino acid and nucleotide sequences encoding one or more particular plant proteins. The skilled artisan, having the benefit of the sequences as reported herein, may now use all or a substantial portion of the disclosed sequences for purposes known to those skilled in this art. Accordingly, the instant invention comprises the complete sequences as reported in the accompanying Sequence Listing, as well as substantial portions of those sequences as defined above.

"Codon degeneracy" refers to divergence in the genetic code permitting variation of the nucleotide sequence without effecting the amino acid sequence of an encoded polypeptide. Accordingly, the instant invention relates to any nucleic acid fragment that encodes all or a substantial portion of the amino acid sequence encoding the acid or the neutral triacylglycerol lipase proteins as set forth in SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32 and 34. The skilled artisan is well aware of the "codon-bias" exhibited by a specific host cell in usage of nucleotide codons to specify a given amino acid. Therefore, when synthesizing a gene for improved expression in a host cell, it is desirable to design the gene such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

"Synthetic genes" can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form gene segments which are then enzymatically assembled to construct the entire gene. "Chemically synthesized", as related to a sequence of DNA, means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of DNA may be accomplished using well established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the genes can be tailored for optimal gene expression based on optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased

towards those codons favored by the host. Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

“Gene” refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. “Native gene” refers to a gene as found in nature with its own regulatory sequences. “Chimeric gene” refers any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. “Endogenous gene” refers to a native gene in its natural location in the genome of an organism. A “foreign” gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A “transgene” is a gene that has been introduced into the genome by a transformation procedure.

“Coding sequence” refers to a DNA sequence that codes for a specific amino acid sequence. “Regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

“Promoter” refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an “enhancer” is a DNA sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters which cause a gene to be expressed in most cell types at most times are commonly referred to as “constitutive promoters”. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg, (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of different lengths may have identical promoter activity.

The "translation leader sequence" refers to a DNA sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency. Examples of translation leader sequences have been described (Turner, R. and Foster, G. D. (1995) *Molecular Biotechnology* 3:225).

The "3' non-coding sequences" refer to DNA sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al. (1989) *Plant Cell* 1:671-680.

"RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into protein by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to RNA transcript that includes the mRNA and so can be translated into protein by the cell. "Antisense RNA" refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (U.S. Patent No. 5,107,065, incorporated herein by reference). The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. "Functional RNA" refers to sense RNA, antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes.

The term "operably linked" refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

The term "expression", as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from the nucleic acid fragment of the invention. Expression may also refer to translation of mRNA into a polypeptide. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein. "Overexpression" refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or

non-transformed organisms. "Co-suppression" refers to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar foreign or endogenous genes (U.S. Patent No. 5,231,020, incorporated herein by reference).

5 "Altered levels" refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

"Mature" protein refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed.

"Precursor" protein refers to the primary product of translation of mRNA; i.e., with pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular
10 localization signals.

A "chloroplast transit peptide" is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the chloroplast or other plastid types present in the cell in which the protein is made. "Chloroplast transit sequence" refers to a nucleotide sequence that encodes a chloroplast transit peptide. A "signal peptide" is an
15 amino acid sequence which is translated in conjunction with a protein and directs the protein to the secretory system (Chrispeels, J. J., (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53). If the protein is to be directed to a vacuole, a vacuolar targeting signal (*supra*) can further be added, or if to the endoplasmic reticulum, an endoplasmic reticulum retention signal (*supra*) may be added. If the protein is to be directed to the nucleus, any signal
20 peptide present should be removed and instead a nuclear localization signal included (Raikhel (1992) *Plant Phys.* 100:1627-1632).

"Transformation" refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" organisms. Examples of
25 methods of plant transformation include Agrobacterium-mediated transformation (De Blaere et al. (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or "gene gun" transformation technology (Klein T. M. et al. (1987) *Nature (London)* 327:70-73; U.S. Patent No. 4,945,050, incorporated herein by reference).

Standard recombinant DNA and molecular cloning techniques used herein are well
30 known in the art and are described more fully in Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989 (hereinafter "Maniatis").

Nucleic acid fragments encoding at least a portion of several triacylglycerol lipases have been isolated and identified by comparison of random plant cDNA sequences to public
35 databases containing nucleotide and protein sequences using the BLAST algorithms well known to those skilled in the art. Table 1 lists the proteins that are described herein, and the designation of the cDNA clones that comprise the nucleic acid fragments encoding these proteins.

TABLE 1
Triacylglycerol Lipases

Enzyme	Clone	Plant
Triacylglycerol Acid Lipase	cen3n.pk0129.e9	Corn
	Contig of:	Corn
	p0075.cslag33r	
	p0126.cnlay46r	
	p0014.ctuty54r	
	p0102.ceral64r	Corn
	p0126.cnlcm37r	Corn
	ncs.pk0013.h1	<i>Catalpa</i>
	rlr72.pk0015.b2	Rice
	rs11n.pk012.h7	Rice
	Contig of:	Soybean
	sdp3c.pk004.n3	
	ssl.pk0022.a1	
Triacylglycerol Neutral Lipase	Contig of:	Soybean
	s1s1c.pk009.o2	
	srr1c.pk001.m19	
	sre.pk0004.d7	
	cr1n.pk0145.c6	Corn
	Contig of:	Corn
	p0010.cbpbe40r	
	p0083.cldcq17r	
	p0048.cqlac25r	
	p0118.chsbw59r	
	cr1.pk0011.c9	
	cdolc.pk002.c22	
	Contig of:	Corn
	p0037.crwan02r	
	p0004.cb1fm22r	
	p0004.cb1ei43r	
	cco1n.pk068.o9	
	p0093.cssao39r	
	cr1n.pk0127.h8	
	rdr1f.pk002.f11	Rice
	Contig of:	Soybean
	sah1c.pk001.k20	
	sre.pk0058.b1	
	sr1.pk0079.e1	Soybean
	wr1.pk0115.f5	Wheat

5 The nucleic acid fragments of the instant invention may be used to isolate cDNAs and genes encoding homologous proteins from the same or other plant species. Isolation of homologous genes using sequence-dependent protocols is well known in the art. Examples of sequence-dependent protocols include, but are not limited to, methods of nucleic acid

hybridization, and methods of DNA and RNA amplification as exemplified by various uses of nucleic acid amplification technologies (e.g., polymerase chain reaction, ligase chain reaction).

For example, genes encoding other acid triacylglycerol lipases, either as cDNAs or
5 genomic DNAs, could be isolated directly by using all or a portion of the instant nucleic acid fragments as DNA hybridization probes to screen libraries from any desired plant employing methodology well known to those skilled in the art. Specific oligonucleotide probes based upon the instant nucleic acid sequences can be designed and synthesized by methods known in the art (Maniatis). Moreover, the entire sequences can be used directly to synthesize
10 DNA probes by methods known to the skilled artisan such as random primer DNA labeling, nick translation, or end-labeling techniques, or RNA probes using available *in vitro* transcription systems. In addition, specific primers can be designed and used to amplify a part or all of the instant sequences. The resulting amplification products can be labeled directly during amplification reactions or labeled after amplification reactions, and used as
15 probes to isolate full length cDNA or genomic fragments under conditions of appropriate stringency.

In addition, two short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols to amplify longer nucleic acid fragments encoding homologous genes from DNA or RNA. The polymerase chain reaction may also be
20 performed on a library of cloned nucleic acid fragments wherein the sequence of one primer is derived from the instant nucleic acid fragments, and the sequence of the other primer takes advantage of the presence of the polyadenylic acid tracts to the 3' end of the mRNA precursor encoding plant genes. Alternatively, the second primer sequence may be based upon sequences derived from the cloning vector. For example, the skilled artisan can follow
25 the RACE protocol (Frohman et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:8998) to generate cDNAs by using PCR to amplify copies of the region between a single point in the transcript and the 3' or 5' end. Primers oriented in the 3' and 5' directions can be designed from the instant sequences. Using commercially available 3' RACE or 5' RACE systems (BRL), specific 3' or 5' cDNA fragments can be isolated (Ohara et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:5673; Loh et al. (1989) *Science* 243:217). Products generated by the 3' and 5' RACE procedures can be combined to generate full-length cDNAs (Frohman, M. A. and Martin, G. R., (1989) *Techniques* 1:165).
30

Availability of the instant nucleotide and deduced amino acid sequences facilitates immunological screening of cDNA expression libraries. Synthetic peptides representing
35 portions of the instant amino acid sequences may be synthesized. These peptides can be used to immunize animals to produce polyclonal or monoclonal antibodies with specificity for peptides or proteins comprising the amino acid sequences. These antibodies can be then be used to screen cDNA expression libraries to isolate full-length cDNA clones of interest (Lerner, R. A. (1984) *Adv. Immunol.* 36:1; Maniatis).

The nucleic acid fragments of the instant invention may be used to create transgenic plants in which the disclosed acid or neutral triacylglycerol lipases are present at higher or lower levels than normal or in cell types or developmental stages in which they are not normally found. This would have the effect of altering the level of triacylglycerol and cholesteryl esters in those cells. Accumulation of fatty acids with unusual structures may be a positive phenotype in plants used for foods. Triacylglycerol lipases may also be useful in processing of plant seed oils and the development of novel seed oils.

Overexpression of the acid or the neutral triacylglycerol lipases of the instant invention may be accomplished by first constructing a chimeric gene in which the coding region is operably linked to a promoter capable of directing expression of a gene in the desired tissues at the desired stage of development. For reasons of convenience, the chimeric gene may comprise promoter sequences and translation leader sequences derived from the same genes. 3' Non-coding sequences encoding transcription termination signals may also be provided. The instant chimeric gene may also comprise one or more introns in order to facilitate gene expression.

Plasmid vectors comprising the instant chimeric gene can then be constructed. The choice of plasmid vector is dependent upon the method that will be used to transform host plants. The skilled artisan is well aware of the genetic elements that must be present on the plasmid vector in order to successfully transform, select and propagate host cells containing the chimeric gene. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al. (1985) *EMBO J.* 4:2411-2418; De Almeida et al. (1989) *Mol. Gen. Genetics* 218:78-86), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

For some applications it may be useful to direct the instant triacylglycerol lipase to different cellular compartments, or to facilitate its secretion from the cell. It is thus envisioned that the chimeric gene described above may be further supplemented by altering the coding sequence to encode a acid triacylglycerol lipase with appropriate intracellular targeting sequences such as transit sequences (Keegstra, K. (1989) *Cell* 56:247-253), signal sequences or sequences encoding endoplasmic reticulum localization (Chrispeels, J. J., (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53), or nuclear localization signals (Raikhel, N. (1992) *Plant Phys.* 100:1627-1632) added and/or with targeting sequences that are already present removed. While the references cited give examples of each of these, the list is not exhaustive and more targeting signals of utility may be discovered in the future.

It may also be desirable to reduce or eliminate expression of genes encoding acid or neutral triacylglycerol lipases in plants for some applications. In order to accomplish this, a chimeric gene designed for co-suppression of the instant triacylglycerol lipase can be

constructed by linking a gene or gene fragment encoding an acid or a neutral triacylglycerol lipase to plant promoter sequences. Alternatively, a chimeric gene designed to express antisense RNA for all or part of the instant nucleic acid fragment can be constructed by linking the gene or gene fragment in reverse orientation to plant promoter sequences. Either
5 the co-suppression or antisense chimeric genes could be introduced into plants via transformation wherein expression of the corresponding endogenous genes are reduced or eliminated.

The instant acid or neutral triacylglycerol lipases (or portions thereof) may be produced in heterologous host cells, particularly in the cells of microbial hosts, and can be
10 used to prepare antibodies to these proteins by methods well known to those skilled in the art. The antibodies are useful for detecting acid or neutral triacylglycerol lipases *in situ* in cells or *in vitro* in cell extracts. Preferred heterologous host cells for production of the instant acid or neutral triacylglycerol lipases are microbial hosts. Microbial expression systems and expression vectors containing regulatory sequences that direct high level
15 expression of foreign proteins are well known to those skilled in the art. Any of these could be used to construct a chimeric gene for production of the instant acid or neutral triacylglycerol lipase. This chimeric gene could then be introduced into appropriate microorganisms via transformation to provide high level expression of the encoded triacylglycerol lipase. An example of a vector for high level expression of the instant acid or
20 neutral triacylglycerol lipase in a bacterial host is provided (Example 7).

All or a substantial portion of the nucleic acid fragments of the instant invention may also be used as probes for genetically and physically mapping the genes that they are a part of, and as markers for traits linked to those genes. Such information may be useful in plant breeding in order to develop lines with desired phenotypes. For example, the instant nucleic
25 acid fragments may be used as restriction fragment length polymorphism (RFLP) markers. Southern blots (Maniatis) of restriction-digested plant genomic DNA may be probed with the nucleic acid fragments of the instant invention. The resulting banding patterns may then be subjected to genetic analyses using computer programs such as MapMaker (Lander et al. (1987) *Genomics* 1:174-181) in order to construct a genetic map. In addition, the nucleic
30 acid fragments of the instant invention may be used to probe Southern blots containing restriction endonuclease-treated genomic DNAs of a set of individuals representing parent and progeny of a defined genetic cross. Segregation of the DNA polymorphisms is noted and used to calculate the position of the instant nucleic acid sequence in the genetic map previously obtained using this population (Botstein, D. et al. (1980) *Am. J. Hum. Genet.*
35 32:314-331).

The production and use of plant gene-derived probes for use in genetic mapping is described in R. Bernatzky, R. and Tanksley, S. D. (1986) *Plant Mol. Biol. Reporter* 4(1):37-41. Numerous publications describe genetic mapping of specific cDNA clones using the methodology outlined above or variations thereof. For example, F2 intercross

populations, backcross populations, randomly mated populations, near isogenic lines, and other sets of individuals may be used for mapping. Such methodologies are well known to those skilled in the art.

5 Nucleic acid probes derived from the instant nucleic acid sequences may also be used for physical mapping (i.e., placement of sequences on physical maps; see Hoheisel, J. D., et al. In: *Nonmammalian Genomic Analysis: A Practical Guide*, Academic press 1996, pp. 319-346, and references cited therein).

10 In another embodiment, nucleic acid probes derived from the instant nucleic acid sequences may be used in direct fluorescence *in situ* hybridization (FISH) mapping (Trask, B. J. (1991) *Trends Genet.* 7:149-154). Although current methods of FISH mapping favor use of large clones (several to several hundred KB; see Laan, M. et al. (1995) *Genome Research* 5:13-20), improvements in sensitivity may allow performance of FISH mapping using shorter probes.

15 A variety of nucleic acid amplification-based methods of genetic and physical mapping may be carried out using the instant nucleic acid sequences. Examples include allele-specific amplification (Kazazian, H. H. (1989) *J. Lab. Clin. Med.* 114(2):95-96), polymorphism of PCR-amplified fragments (CAPS; Sheffield, V. C. et al. (1993) *Genomics* 16:325-332), allele-specific ligation (Landegren, U. et al. (1988) *Science* 241:1077-1080), nucleotide extension reactions (Sokolov, B. P. (1990) *Nucleic Acid Res.* 18:3671), Radiation
20 Hybrid Mapping (Walter, M. A. et al. (1997) *Nature Genetics* 7:22-28) and Happy Mapping (Dear, P. H. and Cook, P. R. (1989) *Nucleic Acid Res.* 17:6795-6807). For these methods, the sequence of a nucleic acid fragment is used to design and produce primer pairs for use in the amplification reaction or in primer extension reactions. The design of such primers is well known to those skilled in the art. In methods employing PCR-based genetic mapping,
25 it may be necessary to identify DNA sequence differences between the parents of the mapping cross in the region corresponding to the instant nucleic acid sequence. This, however, is generally not necessary for mapping methods.

Loss of function mutant phenotypes may be identified for the instant cDNA clones either by targeted gene disruption protocols or by identifying specific mutants for these
30 genes contained in a maize population carrying mutations in all possible genes (Ballinger and Benzer, (1989) *Proc. Natl. Acad. Sci USA* 86:9402; Koes et al. (1995) *Proc. Natl. Acad. Sci USA* 92:8149; Bensen et al. (1995) *Plant Cell* 7:75). The latter approach may be accomplished in two ways. First, short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols in conjunction with a mutation tag sequence
35 primer on DNAs prepared from a population of plants in which Mutator transposons or some other mutation-causing DNA element has been introduced (see Bensen, *supra*). The amplification of a specific DNA fragment with these primers indicates the insertion of the mutation tag element in or near the plant gene encoding the acid or the neutral triacylglycerol lipase. Alternatively, the instant nucleic acid fragment may be used as a

hybridization probe against PCR amplification products generated from the mutation population using the mutation tag sequence primer in conjunction with an arbitrary genomic site primer, such as that for a restriction enzyme site-anchored synthetic adaptor. With either method, a plant containing a mutation in the endogenous gene encoding an acid or a neutral triacylglycerol lipase can be identified and obtained. This mutant plant can then be used to determine or confirm the natural function of the acid or the neutral triacylglycerol lipase gene product.

EXAMPLES

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

EXAMPLE 1

Composition of cDNA Libraries: Isolation and Sequencing of cDNA Clones

cDNA libraries representing mRNAs from various *Catalpa*, corn, rice, soybean and wheat tissues were prepared. The characteristics of the libraries are described below.

TABLE 2

cDNA Libraries from Catalpa, Corn, Rice, Soybean and Wheat

Library	Tissue	Clone
cco1n	Corn Cob of 67 Day Old Plants Grown in Green House*	cco1n.pk068.o9
cdolc	Corn Ovary (including pedicel and glumes), 5 Days After Silking	cdolc.pk002.c22
cen3n	Corn Endosperm 20 Days After Pollination*	cen3n.pk0129.e9
cr1	Corn Root From 7 Day Old Seedlings	cr1.pk0011.c9
cr1n	Corn Root From 7 Day Old Seedlings*	cr1n.pk0127.h8 cr1n.pk0145.c6
ncs	<i>Catalpa speciosa</i> Developing Seed	ncs.pk0013.h1
p0004	Corn Immature Ear	p0004.cb1ei43r
p0010	Corn Log Phase Suspension Cells Treated With A23187**	p0004.cb1fm22r p0010.cbpbe40r
p0014	Corn Leaves 7 and 8 From 3 Foot-Tall Plant	p0014.ctuty54r
p0037	Corn V5 Stage Roots Infested With Corn Root Worm	p0037.crwan02r
p0048	Corn Embryo (Axis and Scutellum) One Day After Germination	p0048.cqlac25r
p0075	Corn Shoot And Leaf Material From Dark-Grown 7 Day-Old Seedlings	p0075.cslag33r

Library	Tissue	Clone
p0083	Corn Whole Kernels 7 Days After Pollination	p0083.cldcq17r
p0093	Corn Stalk And Shank, 2-3 Weeks After Pollen Shed*	p0093.cssao39r
p0102	Corn Early Meiosis Tassels*	p0102.ceral64r
p0118	Corn Stem Tissue Pooled From the 4 to 5 Internodes Subtending The Tassel At Stages V8-V12, Night Harvested*	p0118.chsbw59r
p0126	Corn Leaf Tissue From V8-V10 Stages, Pooled, Night-Harvested	p0126.cnlay46r p0126.cnlcm37r
rdr1f	Rice Developing Root of 10 Day Old Plant	rdr1f.pk002.f11
rlr72	Rice Leaf 15 Days After Germination, 72 Hours After Infection of Strain <i>Magaporthe grisea</i> 4360-R-62 (AVR2-YAMO); Resistant	rlr72.pk0015.b2
rsl1n	Rice 15-Day-Old Seedling*	rsl1n.pk012.h7
sahlc	Soybean Sprayed With Authority™ Herbicide	sahlc.pk001.k20
sdp3c	Soybean Developing Pods (8-9 mm)	sdp3c.pk004.n3
sls1c	Soybean Infected With <i>Sclerotinia sclerotiorum</i> Mycelium	sls1c.pk009.o2
srl	Soybean Root	srl.pk0079.e1
sre	Soybean Root Elongation Zone 4 to 5 Days After Germination	sre.pk0004.d7 sre.pk0058.b1
srr1c	Soybean 8-Day-Old Root	srr1c.pk001.m19
ssl	Soybean Seedling 5-10 Days After Germination	ssl.pk0022.a1
wrl	Wheat Root From 7 Day Old Seedling	wrl.pk0115.f5

*These libraries were normalized essentially as described in U.S. Patent No. 5,482,845

**A23187 is commercially available from several sources including Calbiochem.

- 5 cDNA libraries were prepared in Uni-ZAP™ XR vectors according to the manufacturer's protocol (Stratagene Cloning Systems, La Jolla, CA). Conversion of the Uni-ZAP™ XR libraries into plasmid libraries was accomplished according to the protocol provided by Stratagene. Upon conversion, cDNA inserts were contained in the plasmid vector pBluescript. cDNA inserts from randomly picked bacterial colonies containing
- 10 recombinant pBluescript plasmids were amplified via polymerase chain reaction using primers specific for vector sequences flanking the inserted cDNA sequences or plasmid DNA was prepared from cultured bacterial cells. Amplified insert DNAs or plasmid DNAs were sequenced in dye-primer sequencing reactions to generate partial cDNA sequences (expressed sequence tags or "ESTs"; see Adams, M. D. et al. (1991) *Science* 252:1651).
- 15 The resulting ESTs were analyzed using a Perkin Elmer Model 377 fluorescent sequencer.

EXAMPLE 2

Identification of cDNA Clones

ESTs encoding triacylglycerol lipases were identified by conducting BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al. (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/) searches for similarity to sequences contained in the BLAST "nr" database (comprising all non-redundant GenBank CDS translations, sequences derived from the 3-dimensional structure Brookhaven Protein Data Bank, the last major release of the SWISS-PROT protein sequence database, EMBL, and DDBJ databases). The cDNA sequences obtained in Example 1 were analyzed for similarity to all publicly available DNA sequences contained in the "nr" database using the BLASTN algorithm provided by the National Center for Biotechnology Information (NCBI). The DNA sequences were translated in all reading frames and compared for similarity to all publicly available protein sequences contained in the "nr" database using the BLASTX algorithm (Gish, W. and States, D. J. (1993) *Nature Genetics* 3:266-272) provided by the NCBI. For convenience, the P-value (probability) of observing a match of a cDNA sequence to a sequence contained in the searched databases merely by chance as calculated by BLAST are reported herein as "pLog" values, which represent the negative of the logarithm of the reported P-value. Accordingly, the greater the pLog value, the greater the likelihood that the cDNA sequence and the BLAST "hit" represent homologous proteins.

EXAMPLE 3

Characterization of cDNA Clones Encoding Acid Triacylglycerol Lipases

The BLASTX search using the EST sequences from clones cen3n.pk0129.e9, ncs.pk0013.h1, a contig sequence assembled from the EST sequences from clones rlr72.pk0015.b2 and rr1.pk0051.f10, a contig sequence assembled from the EST sequences of clones ssl.pk0022.a1 and sr1.pk0098.b11, and a contig sequence assembled from the EST sequences from clones sre.pk0004.d7 and sre.pk0001.b2 revealed similarity of the proteins encoded by the cDNAs and the contigs to acid triacylglycerol lipases from human and rat (GenBank Accession Nos. are listed below). The BLAST results for each of these ESTs and contigs are shown in Table 3:

TABLE 3
**BLAST Results for Clones Encoding Polypeptides Homologous
to Acid Triacylglycerol Lipases**

Clone	Organism	GenBank Accession No.	BLAST pLog Score
cen3n.pk0129.e9	Human	X05997	14.52
ncs.pk0013.h1	Rat	X02309	14.70
Contig of rlr72.pk0015.b2 rr1.pk0051.f10	Human	U08464	16.40
Contig of ssl.pk0022.a1 sr1.pk0098.b11	Rat	X02309	15.22
Contig of sre.pk0004.d7 sre.pk0001.b2	Human	X76488	22.00

- 5 TBLASTN analysis of the proprietary plant EST database indicated that other corn, rice and soybean sequences also encoded acid triacylglycerol lipases. The BLASTX search using the contig sequences assembled with the EST sequences from clones p0075.cslag33r, p0126.cnlay46r and p0014.ctuty54r revealed similarity of the proteins encoded by the cDNAs to acid triacylglycerol lipase from *Homo sapiens* (NCBI General Identifier
- 10 No. 505053). The BLASTX search using the EST sequences from clones p0102.ceral64r and using the contig sequences assembled from the entire cDNA insert in clone ssl.pk0022.a1 and the EST sequences from clone sdp3c.pk004.n3 revealed similarity of the proteins encoded by the cDNAs to acid triacylglycerol lipase from *Canis familiaris* (NCBI General Identifier No. 3041702). The BLASTX search using the EST sequences from clone
- 15 p0126.cnlcm37r revealed similarity of the proteins encoded by the cDNAs to *Drosophila melanogaster* (NCBI General Identifier No. 2894442). The BLASTX search using the EST sequences from clone rsl1n.pk012.h7 revealed similarity of the proteins encoded by the cDNAs to acid triacylglycerol lipase from *Rattus norvegicus* (NCBI General Identifier No. 126307). The BLAST results for each of these sequences is shown in Table 4:

20

TABLE 4

BLAST Results for Clones Encoding Polypeptides Homologous
to Acid Triacylglycerol Lipase

Clone	NCBI General Identifier No.	BLAST pLog Score
Contig of: p0075.cslag33r p0126.cnlay46r p0014.ctuty54r	505053	35.00
p0102.ceral64r	3041702	11.30
p0126.cnlcm37r	2894442	10.40
rsl1n.pk012.h7	126307	7.00

- 5 The sequence of the entire cDNA insert in clone cen3n.pk0129.e9 was determined and is shown in SEQ ID NO:1; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:2. The amino acid sequence set forth in SEQ ID NO:2 was evaluated by BLASTP, yielding a pLog value of 15.00 versus the *Homo sapiens* sequence (NCBI General Identifier No. 126306). The sequence of the 3'-terminal portion from clone ncs.pk0013.h1 is shown in
- 10 SEQ ID NO:3; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:4. The sequence of the 5'-terminal portion from clone ncs.pk0013.h1 is shown in SEQ ID NO:5; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:6. The sequence of the contig assembled from the EST sequences from clones p0075.cslag33r, p0126.cnlay46r and p0014.ctuty54r is shown in SEQ ID NO:7, the deduced amino acid
- 15 sequence of this cDNA is shown in SEQ ID NO:8. The sequence of a portion of the cDNA insert from clone p0102.ceral64r is shown in SEQ ID NO:9; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:10. The sequence of a portion of the cDNA insert from clone p0126.cnlcm37r is shown in SEQ ID NO:11; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:12. The sequence of the entire cDNA insert
- 20 in clone rlr72.pk0015.b2 was determined and is shown in SEQ ID NO:13; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:14. The amino acid sequence set forth in SEQ ID NO:14 was evaluated by BLASTP, yielding a pLog value of 53.30 versus the *C. elegans* sequence (NCBI General Identifier No. 3165581). The sequence of a portion of the cDNA insert from clone rsl1n.pk012.h7 is shown in SEQ ID NO:15; the
- 25 deduced amino acid sequence of this cDNA is shown in SEQ ID NO:16. The sequence of the entire cDNA insert in clone ssl.pk0022.a1 was determined and a contig assembled with this sequence and the EST sequences from clone sdp3c.pk004.n3. The sequence of this contig is shown in SEQ ID NO:17; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:18. The amino acid sequence set forth in SEQ ID NO:18 was evaluated by
- 30 BLASTP, yielding a pLog value of 59.40 versus the *C. familiaris* sequence (NCBI General Identifier No. 3041702). The sequence of the entire cDNA insert in clone sre.pk0004.d7 was determined and a contig assembled with this sequence and the EST sequences from

clones sls1c.pk009.o2 and srr1c.pk001.m19. The sequence of this contig is shown in SEQ ID NO:19; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:20. The amino acid sequence set forth in SEQ ID NO:20 was evaluated by BLASTP, yielding a pLog value of 48.70 versus the *C. elegans* sequence (NCBI General Identifier No. 3165581).

5 Figure 1 presents an alignment of the amino acid sequences set forth in SEQ ID NOs:14, 18 and 20 with the *Canis familiaris* sequence (NCBI General Identifier No. 3041702; SEQ ID NO:35) and the *Caenorhabditis elegans* sequence (NCBI General Identifier No. 3165581; SEQ ID NO:36). The data in Table 5 presents a calculation of the percent similarity of the amino acid sequences set forth in SEQ ID NOs:2, 4, 6, 8, 10, 12, 14,
10 18 and 20 and the *Caenorhabditis elegans* sequence.

15 **TABLE 5**
Percent Similarity of Amino Acid Sequences Deduced From the Nucleotide Sequences of cDNA Clones Encoding Polypeptides Homologous to Acid Triacylglycerol Lipase

Clone	SEQ ID NO.	Percent Identity to	
		3041702	3165581
cen3n.pk0129.e9: fis	2	27.1	22.9
ncs.pk0013.h1.fis1	4	27.4	21.4
ncs.pk0013.h1.fis2	6	30.6	29.9
p0075.cslag33r	8	22.0	23.1
p0126.cnlay46r			
p0014.ctuty54r			
p0102.ceral64r	10	28.8	22.4
p0126.cnlcm37r	12	26.7	22.2
rlr72.pk0015.b2: fis	14	24.9	25.6
rsl1n.pk012.h7	16	22.5	22.5
sdp3c.pk004.n3	18	27.4	23.1
ssl.pk0022.a1.fis1			
sls1c.pk009.o2	20	23.1	24.8
srr1c.pk001.m19			
sre.pk0004.d7.fis1			

Sequence alignments and percent similarity calculations were performed using the Megalign program of the LASARGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the amino acid sequences was performed using the Clustal
20 method of alignment (Higgins, D. G. and Sharp, P. M. (1989) *CABIOS* 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10).

Sequence alignments and BLAST scores and probabilities indicate that the instant nucleic acid fragments encode an entire rice acid triacylglycerol lipase, two different entire soybean acid triacylglycerol lipases, portions from several different corn acid triacylglycerol
25 lipases, portions of a *Catalpa* acid triacylglycerol lipase and a portion of a rice acid

triacylglycerol lipase. These sequences represent the first plant sequences encoding acid triacylglycerol lipases.

EXAMPLE 4

Characterization of cDNA Clones Encoding Neutral Triacylglycerol Lipases

5 The BLASTX search using the contig sequence assembled from the EST sequences from clones cr1n.pk0127.h8 and cr1n.pk0134.d3, and EST sequences from clones cr1n.pk0145.c6, sl.03b01, se3.01a04, sfl1.pk0049.d11, sr1.pk0079.e1, sr1.pk0030.g8, sre.pk0058.b1, wlln.pk0014.e10, wlln.pk0038.e3 and wr1.pk0115.f5 revealed similarity of the proteins encoded by the contig and the cDNAs to neutral triacylglycerol lipases from
10 several organisms. Table 6 shows the BLAST results for the contig and each of the ESTs, the NCBI database accession number, and the organism the closest art sequence is derived from:

TABLE 6

15 BLAST Results for Clones Encoding Polypeptides Homologous to Neutral Triacylglycerol Lipases

Clone	Organism	NCBI Accession No.	BLAST pLog Score
Contig of: cr1n.pk0127.h8 cr1n.pk0134.d3	<i>Thermomyces lanuginosus</i>	999873	10.00
cr1n.pk0145.c6	<i>Caenorhabditis elegans</i>	927399	8.70
sr1.pk0079.e1	<i>Rhizopus niveus</i>	251079	6.70
sre.pk0058.b1	<i>Rhizomucor miehei</i>	417256	8.10
wr1.pk0115.f5	<i>Rhizomucor miehei</i>	82777	6.00

TBLASTN analysis of the proprietary plant EST database indicated that rice clones as well as other corn and soybean clones also encode neutral triacylglycerol lipases. The
20 BLASTX search using the contig sequences assembled from clones p0010.cbpb40r, p0083.cldcql7r, p0048.cqlac25r, p0118.chsbw59r, cr1.pk0011.c9 and cdo1c.pk002.c22 and using the EST sequences from clone rdr1f.pk002.f11 revealed similarity of the proteins encoded by the cDNAs to neutral triacylglycerol lipase from *C. elegans* (NCBI General Identifier No.3877256). The BLAST results for each of these sequences are shown in
25 Table 7:

TABLE 7

BLAST Results for Clones Encoding Polypeptides Homologous
to Neutral Triacylglycerol Lipases

Clone	Organism	NCBI General Identifier No.	BLAST pLog Score
crln.pk0145.c6	<i>Caenorhabditis elegans</i>	3877256	9.30
Contig of: p0010.cbpbe40r p0083.cldcq17r p0048.cqlac25r p0118.chsbw59r cr1.pk0011.c9 cdolc.pk002.c22	<i>Caenorhabditis elegans</i>	3877256	18.40
Contig of: p0037.crwan02r p0004.cb1fm22r p0004.cb1ei43r cco1n.pk068.o9 p0093.cssao39r crln.pk0127.h8	<i>Thermomyces lanuginosus</i>	2997733	6.15
rdr1f.pk002.f11	<i>Caenorhabditis elegans</i>	3877256	10.22
Contig of: sah1c.pk001.k20 sre.pk0058.b1	<i>Rhizomucor miehei</i>	417256	6.22
srl.pk0079.e1	<i>Rhizopus niveus</i>	3299795	5.70
wrl.pk0115.f5	<i>Caenorhabditis elegans</i>	3877256	14.00

- 5 The sequence of the entire cDNA insert in clone crln.pk0145.c6 was determined and is shown in SEQ ID NO:21; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:22. The amino acid sequence set forth in SEQ ID NO:2 was evaluated by BLASTP, yielding a pLog value of 10.70 versus the *C. elegans* sequence. The sequence of the contig assembled from a portion of the cDNA insert in clones p0010.cbpbe40r,
- 10 p0083.cldcq17r, p0048.cqlac25r, p0118.chsbw59r, cr1.pk0011.c9 and cdolc.pk002.c22 is shown in SEQ ID NO:23; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:23. The sequence of the entire cDNA insert in clone crln.pk0127.h8 was determined and a contig assembled with this sequence and the sequence from a portion of the cDNA insert in clones p0037.crwan02r, p0004.cb1fm22r, p0004.cb1ei43r, cco1n.pk068.o9 and
- 15 p0093.cssao39r. The sequence of this contig is shown in SEQ ID NO:25; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:26. The amino acid sequence set forth in SEQ ID NO:4 was evaluated by BLASTP, yielding a pLog value of 9.70 versus the *Thermomyces lanuginosus* sequence. The sequence of a portion of the cDNA insert from clone rdr1f.pk002.f11 is shown in SEQ ID NO:27; the deduced amino acid sequence of
- 20 this cDNA is shown in SEQ ID NO:28. The sequence of the entire cDNA insert in clone sre.pk0058.b1 was determined and a contig assembled with this sequence and the sequence

of a portion of the cDNA insert in clone sah1c.pk001.k20. The sequence of this contig is shown in SEQ ID NO:29; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:30. The amino acid sequence set forth in SEQ ID NO:30 was evaluated by BLASTP, yielding a pLog value of 8.05 versus the *Rhizomucor miehei* sequence. The sequence of the entire cDNA insert in clone srl.pk0079.e1 was determined and is shown in SEQ ID NO:31; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:32. The amino acid sequence set forth in SEQ ID NO:32 was evaluated by BLASTP, yielding a pLog value of 7.52 versus the *Rhizopus niveus* sequence. The sequence of the entire cDNA insert in clone wr1.pk0115.f5 was determined and is shown in SEQ ID NO:33; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:34. The amino acid sequence set forth in SEQ ID NO:34 was evaluated by BLASTP, yielding a pLog value of 13.52 versus the *Caenorhabditis elegans* sequence.

The data in Table 8 presents a calculation of the percent similarity of the amino acid sequences set forth in SEQ ID NOs:22, 24, 26, 28, 30, 32 and 34 and the *Caenorhabditis elegans*, *Rhizomucor miehei* and *Thermomyces lanuginosus* sequences.

TABLE 8

Percent Similarity of Amino Acid Sequences Deduced From the Nucleotide Sequences of cDNA Clones Encoding Polypeptides Homologous to Neutral Triacylglycerol Lipase

Clone	SEQ ID NO.	Percent Similarity to		
		3877256	2997733	417256
cr1n.pk0145.c6	22	15.1	13.2	16.8
Contig of:	24	60.5	17.5	22.9
p0010.cbpbe40r				
p0083.cldcq17r				
p0048.cqlac25r				
p0118.chsbw59r				
cr1.pk0011.c9				
cdolc.pk002.c22				
Contig of:	26	18.5	14.3	15.1
p0037.crwan02r				
p0004.cb1fm22r				
p0004.cb1ei43r				
cco1n.pk068.o9				
p0093.cssao39r				
cr1n.pk0127.h8				
rdr1f.pk002.f11	28	12.6	20.6	22.9
Contig of:	32	15.1	10.5	17.0
sah1c.pk001.k20				
sre.pk0058.b1				
srl.pk0079.e1	34	14.3	21.1	24.6
wr1.pk0115.f5	36	37.0	22.0	26.0

Sequence alignments and percent similarity calculations were performed using the Megalign program of the LASARGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the amino acid sequences was performed using the Clustal method of alignment (Higgins, D. G. and Sharp, P. M. (1989) *CABIOS*, 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10).

Sequence alignments and BLAST scores and probabilities indicate that the instant nucleic acid fragments encode three different corn neutral triacylglycerol lipases (one portion and two entire or nearly entire), two different soybean triacylglycerol lipases (one portion and one nearly entire) and a portion of a wheat triacylglycerol lipase. These sequences represent the first monocot and soybean sequences encoding neutral triacylglycerol lipases.

EXAMPLE 5

Expression of Chimeric Genes in Monocot Cells

A chimeric gene comprising a cDNA encoding triacylglycerol lipases in sense orientation with respect to the maize 27 kD zein promoter that is located 5' to the cDNA fragment, and the 10 kD zein 3' end that is located 3' to the cDNA fragment, can be constructed. The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites (Nco I or Sma I) can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the digested vector pML103 as described below. Amplification is then performed in a standard PCR. The amplified DNA is then digested with restriction enzymes Nco I and Sma I and fractionated on an agarose gel. The appropriate band can be isolated from the gel and combined with a 4.9 kb Nco I-Sma I fragment of the plasmid pML103. Plasmid pML103 has been deposited under the terms of the Budapest Treaty at ATCC (American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209), and bears accession number ATCC 97366. The DNA segment from pML103 contains a 1.05 kb Sal I-Nco I promoter fragment of the maize 27 kD zein gene and a 0.96 kb Sma I-Sal I fragment from the 3' end of the maize 10 kD zein gene in the vector pGem9Zf(+) (Promega). Vector and insert DNA can be ligated at 15°C overnight, essentially as described (Maniatis). The ligated DNA may then be used to transform *E. coli* XL1-Blue (Epicurian Coli XL-1 Blue™; Stratagene). Bacterial transformants can be screened by restriction enzyme digestion of plasmid DNA and limited nucleotide sequence analysis using the dideoxy chain termination method (Sequenase™ DNA Sequencing Kit; U.S. Biochemical). The resulting plasmid construct would comprise a chimeric gene encoding, in the 5' to 3' direction, the maize 27 kD zein promoter, a cDNA fragment encoding a triacylglycerol lipase, and the 10 kD zein 3' region.

The chimeric gene described above can then be introduced into corn cells by the following procedure. Immature corn embryos can be dissected from developing caryopses derived from crosses of the inbred corn lines H99 and LH132. The embryos are isolated 10 to 11 days after pollination when they are 1.0 to 1.5 mm long. The embryos are then placed

with the axis-side facing down and in contact with agarose-solidified N6 medium (Chu et al. (1975) *Sci. Sin. Peking* 18:659-668). The embryos are kept in the dark at 27°C. Friable embryogenic callus consisting of undifferentiated masses of cells with somatic proembryoids and embryoids borne on suspensor structures proliferates from the scutellum of these immature embryos. The embryogenic callus isolated from the primary explant can be cultured on N6 medium and sub-cultured on this medium every 2 to 3 weeks.

The plasmid, p35S/Ac (obtained from Dr. Peter Eckes, Hoechst Ag, Frankfurt, Germany) may be used in transformation experiments in order to provide for a selectable marker. This plasmid contains the *Pat* gene (see European Patent Publication 0 242 236) which encodes phosphinothricin acetyl transferase (PAT). The enzyme PAT confers resistance to herbicidal glutamine synthetase inhibitors such as phosphinothricin. The *pat* gene in p35S/Ac is under the control of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*.

The particle bombardment method (Klein, T. M. et al. (1987) *Nature* 327:70-73) may be used to transfer genes to the callus culture cells. According to this method, gold particles (1 µm in diameter) are coated with DNA using the following technique. Ten µg of plasmid DNAs are added to 50 µL of a suspension of gold particles (60 mg per mL). Calcium chloride (50 µL of a 2.5 M solution) and spermidine free base (20 µL of a 1.0 M solution) are added to the particles. The suspension is vortexed during the addition of these solutions. After 10 minutes, the tubes are briefly centrifuged (5 sec at 15,000 rpm) and the supernatant removed. The particles are resuspended in 200 µL of absolute ethanol, centrifuged again and the supernatant removed. The ethanol rinse is performed again and the particles resuspended in a final volume of 30 µL of ethanol. An aliquot (5 µL) of the DNA-coated gold particles can be placed in the center of a Kapton™ flying disc (Bio-Rad Labs). The particles are then accelerated into the corn tissue with a Biolistic™ PDS-1000/He (Bio-Rad Instruments, Hercules CA), using a helium pressure of 1000 psi, a gap distance of 0.5 cm and a flying distance of 1.0 cm.

For bombardment, the embryogenic tissue is placed on filter paper over agarose-solidified N6 medium. The tissue is arranged as a thin lawn and covered a circular area of about 5 cm in diameter. The petri dish containing the tissue can be placed in the chamber of the PDS-1000/He approximately 8 cm from the stopping screen. The air in the chamber is then evacuated to a vacuum of 28 inches of Hg. The macrocarrier is accelerated with a helium shock wave using a rupture membrane that bursts when the He pressure in the shock tube reaches 1000 psi.

Seven days after bombardment the tissue can be transferred to N6 medium that contains gluphosinate (2 mg per liter) and lacks casein or proline. The tissue continues to grow slowly on this medium. After an additional 2 weeks the tissue can be transferred to fresh N6 medium containing gluphosinate. After 6 weeks, areas of about 1 cm in diameter

of actively growing callus can be identified on some of the plates containing the glufosinate-supplemented medium. These calli may continue to grow when sub-cultured on the selective medium.

Plants can be regenerated from the transgenic callus by first transferring clusters of tissue to N6 medium supplemented with 0.2 mg per liter of 2,4-D. After two weeks the tissue can be transferred to regeneration medium (Fromm et al. (1990) *Bio/Technology* 8:833-839).

EXAMPLE 6

Expression of Chimeric Genes in Dicot Cells

A seed-specific expression cassette composed of the promoter and transcription terminator from the gene encoding the β subunit of the seed storage protein phaseolin from the bean *Phaseolus vulgaris* (Doyle et al. (1986) *J. Biol. Chem.* 261:9228-9238) can be used for expression of the instant triacylglycerol lipase in transformed soybean. The phaseolin cassette includes about 500 nucleotides upstream (5') from the translation initiation codon and about 1650 nucleotides downstream (3') from the translation stop codon of phaseolin. Between the 5' and 3' regions are the unique restriction endonuclease sites Nco I (which includes the ATG translation initiation codon), Sma I, Kpn I and Xba I. The entire cassette is flanked by Hind III sites.

The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the expression vector. Amplification is then performed as described above, and the isolated fragment is inserted into a pUC18 vector carrying the seed expression cassette.

Soybean embryos may then be transformed with the expression vector comprising sequences encoding a triacylglycerol lipase. To induce somatic embryos, cotyledons, 3-5 mm in length dissected from surface sterilized, immature seeds of the soybean cultivar A2872, can be cultured in the light or dark at 26°C on an appropriate agar medium for 6-10 weeks. Somatic embryos which produce secondary embryos are then excised and placed into a suitable liquid medium. After repeated selection for clusters of somatic embryos which multiplied as early, globular staged embryos, the suspensions are maintained as described below.

Soybean embryogenic suspension cultures can be maintained in 35 mL liquid media on a rotary shaker, 150 rpm, at 26°C with florescent lights on a 16:8 hour day/night schedule. Cultures are subcultured every two weeks by inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

Soybean embryogenic suspension cultures may then be transformed by the method of particle gun bombardment (Klein T. M. et al. (1987) *Nature* (London) 327:70-73, U.S.

Patent No. 4,945,050). A DuPont Biolistic™ PDS1000/HE instrument (helium retrofit) can be used for these transformations.

A selectable marker gene which can be used to facilitate soybean transformation is a chimeric gene composed of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812), the hygromycin phosphotransferase gene from plasmid pJR225 (from *E. coli*; Gritz et al. (1983) *Gene* 25:179-188) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*. The seed expression cassette comprising the phaseolin 5' region, the fragment encoding the triacylglycerol lipase and the phaseolin 3' region can be isolated as a restriction fragment. This fragment can then be inserted into a unique restriction site of the vector carrying the marker gene.

To 50 µL of a 60 mg/mL 1 µm gold particle suspension is added (in order): 5 µL DNA (1 µg/µL), 20 µl spermidine (0.1 M), and 50 µL CaCl₂ (2.5 M). The particle preparation is then agitated for three minutes, spun in a microfuge for 10 seconds and the supernatant removed. The DNA-coated particles are then washed once in 400 µL 70% ethanol and resuspended in 40 µL of anhydrous ethanol. The DNA/particle suspension can be sonicated three times for one second each. Five µL of the DNA-coated gold particles are then loaded on each macro carrier disk.

Approximately 300-400 mg of a two-week-old suspension culture is placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue are normally bombarded. Membrane rupture pressure is set at 1100 psi and the chamber is evacuated to a vacuum of 28 inches mercury. The tissue is placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue can be divided in half and placed back into liquid and cultured as described above.

Five to seven days post bombardment, the liquid media may be exchanged with fresh media, and eleven to twelve days post bombardment with fresh media containing 50 mg/mL hygromycin. This selective media can be refreshed weekly. Seven to eight weeks post bombardment, green, transformed tissue may be observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue is removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Each new line may be treated as an independent transformation event. These suspensions can then be subcultured and maintained as clusters of immature embryos or regenerated into whole plants by maturation and germination of individual somatic embryos.

EXAMPLE 7

Expression of Chimeric Genes in Microbial Cells

The cDNAs encoding the instant triacylglycerol lipases can be inserted into the T7 *E. coli* expression vector pBT430. This vector is a derivative of pET-3a (Rosenberg et al. (1987) *Gene* 56:125-135) which employs the bacteriophage T7 RNA polymerase/T7 promoter system. Plasmid pBT430 was constructed by first destroying the EcoR I and

Hind III sites in pET-3a at their original positions. An oligonucleotide adaptor containing EcoR I and Hind III sites was inserted at the BamH I site of pET-3a. This created pET-3aM with additional unique cloning sites for insertion of genes into the expression vector. Then, the Nde I site at the position of translation initiation was converted to an Nco I site using
5 oligonucleotide-directed mutagenesis. The DNA sequence of pET-3aM in this region, 5'-CATATGG, was converted to 5'-CCCATGG in pBT430.

Plasmid DNA containing a cDNA may be appropriately digested to release a nucleic acid fragment encoding the protein. This fragment may then be purified on a 1% NuSieve GTG™ low melting agarose gel (FMC). Buffer and agarose contain 10 µg/ml ethidium
10 bromide for visualization of the DNA fragment. The fragment can then be purified from the agarose gel by digestion with GELase™ (Epicentre Technologies) according to the manufacturer's instructions, ethanol precipitated, dried and resuspended in 20 µL of water. Appropriate oligonucleotide adapters may be ligated to the fragment using T4 DNA ligase (New England Biolabs, Beverly, MA). The fragment containing the ligated adapters can be
15 purified from the excess adapters using low melting agarose as described above. The vector pBT430 is digested, dephosphorylated with alkaline phosphatase (NEB) and deproteinized with phenol/chloroform as described above. The prepared vector pBT430 and fragment can then be ligated at 16°C for 15 hours followed by transformation into DH5 electrocompetent cells (GIBCO BRL). Transformants can be selected on agar plates containing LB media and
20 100 µg/mL ampicillin. Transformants containing the gene encoding the triacylglycerol lipase are then screened for the correct orientation with respect to the T7 promoter by restriction enzyme analysis.

For high level expression, a plasmid clone with the cDNA insert in the correct orientation relative to the T7 promoter can be transformed into *E. coli* strain BL21(DE3)
25 (Studier et al. (1986) *J. Mol. Biol.* 189:113-130). Cultures are grown in LB medium containing ampicillin (100 mg/L) at 25°C. At an optical density at 600 nm of approximately 1, IPTG (isopropylthio-β-galactoside, the inducer) can be added to a final concentration of 0.4 mM and incubation can be continued for 3 h at 25°. Cells are then harvested by centrifugation and re-suspended in 50 µL of 50 mM Tris-HCl at pH 8.0 containing 0.1 mM
30 DTT and 0.2 mM phenyl methylsulfonyl fluoride. A small amount of 1 mm glass beads can be added and the mixture sonicated 3 times for about 5 seconds each time with a microprobe sonicator. The mixture is centrifuged and the protein concentration of the supernatant determined. One µg of protein from the soluble fraction of the culture can be separated by SDS-polyacrylamide gel electrophoresis. Gels can be observed for protein bands migrating
35 at the expected molecular weight.

CLAIMS

What is claimed is:

1. An isolated nucleic acid fragment encoding all or a substantial portion of an acid triacylglycerol lipase comprising a member selected from the group consisting of:
 - 5 (a) an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in a member selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18 and SEQ ID NO:20;
 - 10 (b) an isolated nucleic acid fragment that is substantially similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in a member selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18 and SEQ ID NO:20; and
 - 15 (c) an isolated nucleic acid fragment that is complementary to (a) or (b).
2. The isolated nucleic acid fragment of Claim 1 wherein the nucleotide sequence of the fragment comprises all or a portion of the sequence set forth in a member selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7,
 - 20 SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:17 and SEQ ID NO:19.
3. A chimeric gene comprising the nucleic acid fragment of Claim 1 operably linked to suitable regulatory sequences.
4. A transformed host cell comprising the chimeric gene of Claim 3.
- 25 5. An acid triacylglycerol lipase polypeptide comprising all or a substantial portion of the amino acid sequence set forth in a member selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18 and SEQ ID NO:20.
6. An isolated nucleic acid fragment encoding all or a substantial portion of a
 - 30 neutral triacylglycerol lipase comprising a member selected from the group consisting of:
 - (a) an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in a member selected from the group consisting of SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32 and SEQ ID NO:34;
 - 35 (b) an isolated nucleic acid fragment that is substantially similar to an isolated nucleic acid fragment encoding all or a substantial portion of the amino acid sequence set forth in a member selected from the group consisting of SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32 and SEQ ID NO:34; and

(c) an isolated nucleic acid fragment that is complementary to (a) or (b).

7. The isolated nucleic acid fragment of Claim 6 wherein the nucleotide sequence of the fragment comprises all or a portion of the sequence set forth in a member selected from the group consisting of SEQ ID NO:21, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:31 and SEQ ID NO:33.

8. A chimeric gene comprising the nucleic acid fragment of Claim 6 operably linked to suitable regulatory sequences.

9. A transformed host cell comprising the chimeric gene of Claim 8.

10. A neutral triacylglycerol lipase polypeptide comprising all or a substantial portion of the amino acid sequence set forth in a member selected from the group consisting of SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32 and SEQ ID NO:34.

11. A method of altering the level of expression of a triacylglycerol lipase in a host cell comprising:

15 (a) transforming a host cell with the chimeric gene of any of Claims 3 and 8; and

(b) growing the transformed host cell produced in step (a) under conditions that are suitable for expression of the chimeric gene

wherein expression of the chimeric gene results in production of altered levels of a triacylglycerol lipase in the transformed host cell.

12. A method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a triacylglycerol lipase comprising:

(a) probing a cDNA or genomic library with the nucleic acid fragment of any of Claims 1 and 6;

25 (b) identifying a DNA clone that hybridizes with the nucleic acid fragment of any of Claims 1 and 6;

(c) isolating the DNA clone identified in step (b); and

(d) sequencing the cDNA or genomic fragment that comprises the clone isolated in step (c)

30 wherein the sequenced nucleic acid fragment encodes all or a substantial portion of the amino acid sequence encoding a triacylglycerol lipase.

13. A method of obtaining a nucleic acid fragment encoding a substantial portion of an amino acid sequence encoding a triacylglycerol lipase comprising:

35 (a) synthesizing an oligonucleotide primer corresponding to a portion of the sequence set forth in any of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31 and 33; and

(b) amplifying a cDNA insert present in a cloning vector using the oligonucleotide primer of step (a) and a primer representing sequences of the cloning vector

wherein the amplified nucleic acid fragment encodes a substantial portion of an amino acid sequence encoding a triacylglycerol lipase.

14. The product of the method of Claim 12.
15. The product of the method of Claim 13.

5

FIGURE 1

[illegible]

FIGURE 1 - CONTINUED

```

+   ***   **   ++   +   +++*   *   ++
SEQ ID NO: 37  LHYVGHSQGTIGFIAFSTNP-KLAKRIKTFYALAPVATVKYTETLLNKLMLVPSFLFKL
SEQ ID NO: 38  VYYMGHSQGTITMFSLSKDDGSAKIKKFFALAPIGSVKHIKGFLS---FFANY-FSL
SEQ ID NO: 14  ILYVGHSQGTIMGLAALTM--PEIVKMISAAALLCPISYLDHVSASFV-LRAVAMHLDQM
SEQ ID NO: 18  INYVGHSGLTLVALASFSE--GKLVTLQLKSAALLSPIAYLSHMNTALG-VVAPKSFVGEI
SEQ ID NO: 20  IFVVGHSQGTIISLAFTQ--PEIVEKVEAAALLSPISYLDHVSAPLV-LRMVKMHIDEM
181
240

```

```

*   *   +   ++*   *   *   **   ++   +   *
SEQ ID NO: 37  -----IFGNKIFYPHHFFDQFLATEVCSRETVDL-LCSNALFIICGFDTMNLNMSRLD
SEQ ID NO: 38  EFDGWEDIFGAGEFLPNNWAMKLAAKDICGGLKVEADLCDNVLFLLIAGPESDQWNQTRVP
SEQ ID NO: 14  LV-----TMGIHQLNFRSDMGVQIVDSLCDGEHVD---CNNLLSAITGENC-CFNTSRID
SEQ ID NO: 18  TT-----LFGLAEFNPKGLAVDAFLKSLCAHPGID---CYDLLLTALTGKNC-CLNSSTVD
SEQ ID NO: 20  IL-----TMGIHQLNFKSEWGASLLVSLCDT-RLS---CNDMLSSITGKNC-CFNESRVE
241
300

```

```

++   *   +   *   *   ++   *   +   *   +   +   ++
SEQ ID NO: 37  VYLSHNPAGTSVQNVLHWSQAVKSGKFQAFDWGSPVQNMHHYHQSMPPYYNLTDMHVPPIA
SEQ ID NO: 38  VYATHDPAGTSTQNIIVHWMQMVHHGGVPAYDWGKTNNKKK-YGQANPPEYDFTAIGTKI
SEQ ID NO: 14  YYLEYEPHPSSTKNLHHLFQMIRKGTFAKYDYGLLG-NLRRYGHRLRPPAFDLSSIPESLP
SEQ ID NO: 18  LFLMNEPQSTSTKNMVHLAQTVRLGALTGFENYVRPDYNIMHYGEIFPPIYNLSNIPHDLP
SEQ ID NO: 20  FYLEQEPHPSSTKNLHHLFQMIRKGTYSKYDYGKLG-NLIEYGKFNPPKFDLSRIPKSLP
301
360

```

FIGURE 1 - CONTINUED

SEQ ID NO: 37	V--WNGGNDLLADPHDVDLLLSKL-PNLIYHRK---IPPNHLDFIWMADAPQAVYNEIV
SEQ ID NO: 38	YLYWSDADWLADTPDPDYLLTRLNPAAVAQN--HLPDYNHLDEFTWGLRAPDDIYRPAI
SEQ ID NO: 14	IWMGYGGDLALADVTDVQRTIREL-----GSTPELLYIGDYGHIDFVM SVKAKDDVYVDLI
SEQ ID NO: 18	LFISYGGRDALS DVRDVENLLDKLKFDENKRSVQFIQEYAHADYIMGFNAKD LVYNAVIL
SEQ ID NO: 20	LWMAYGGNDALADITDFQH TLKEL----PSPPEVVYLENYGHVDFILSLQAKQDLYDPMI
	361
	420

SEQ	ID	NO: 37	SMMGTD----	NK
SEQ	ID	NO: 38	KLCTDD--	YL GK
SEQ	ID	NO: 14	RFLRENGW	HSY
SEQ	ID	NO: 18	SFFNHQ----	V
SEQ	ID	NO: 20	SFFKSSGKF	SSM
			421	433

SEQUENCE LISTING

<110> E. I. DU PONT DE NEMOURS AND COMPANY

<120> TRIACYLGLYCEROL LIPASES

<130> BB-1168

<140>

<141>

<150> 60/083,688

<151> 1988-04-30

<160> 36

<170> Microsoft Office 97

<210> 1

<211> 751

<212> DNA

<213> Zea mays

<400> 1

```

gcacgagatc accggcaaga actactgcct caacagctcc gccgtcgacg tcttccctcaa 60
gtacgagccc cagccgacct ccacaaaaac catggtccac ttcgctcaaa ccgtgcgcga 120
cggcgtgctg accaagtacg actacgtgct gccggagcgg aacatcgcca gctacggcca 180
ggccgagccg ccggtgtacc ggatgtccgg catcccgccg agcttcccgc tcttccctcag 240
ctacggcgcc cgggactcgc tcgccgaccc cgccgacgtg cgctcctcc tgcaggacct 300
ccggggccac gaccaggaca agctcacggt gcagtacctg gacaagtctg cgcacctcga 360
cttcatcatc ggcgtctgcg ccaaggacta cgtctacaag gacatgatcg acttccctaaa 420
ccgcttcaac tagtactagc atatataatt gcttcaatcg gtgtcgtctt cagccccagc 480
aggattagac aaaaaaagg ggggacactg cagctcgtaa acgttgtcca tacagattat 540
cagagggtgaa aaccatacat gatgtaattt agcattagat agttaaaaca tggagctgcc 600
tcagtatgga ggattgtcaa ctactctcca tcacagcagt aggtgtgatg tagaagagtg 660
attgtcacac tgtgtgtgtt gcaaatttgc aacacagtga ttactaatat aaaaaaatact 720
cttgagttaa aaaaaaaaaa aaaaaaaaaa a

```

751

<210> 2

<211> 143

<212> PRT

<213> Zea mays

<400> 2

```

His Glu Ile Thr Gly Lys Asn Tyr Cys Leu Asn Ser Ser Ala Val Asp
  1             5             10             15

```

```

Val Phe Leu Lys Tyr Glu Pro Gln Pro Thr Ser Thr Lys Thr Met Val
      20             25             30

```

```

His Phe Ala Gln Thr Val Arg Asp Gly Val Leu Thr Lys Tyr Asp Tyr
      35             40             45

```

```

Val Leu Pro Glu Arg Asn Ile Ala Ser Tyr Gly Gln Ala Glu Pro Pro
      50             55             60

```

```

Val Tyr Arg Met Ser Gly Ile Pro Pro Ser Phe Pro Leu Phe Leu Ser
      65             70             75             80

```

```

Tyr Gly Gly Arg Asp Ser Leu Ala Asp Pro Ala Asp Val Arg Leu Leu
      85             90             95

```

Leu Gln Asp Leu Arg Gly His Asp Gln Asp Lys Leu Thr Val Gln Tyr
 100 105 110

Leu Asp Lys Phe Ala His Leu Asp Phe Ile Ile Gly Val Cys Ala Lys
 115 120 125

Asp Tyr Val Tyr Lys Asp Met Ile Asp Phe Leu Asn Arg Phe Asn
 130 135 140

<210> 3
 <211> 647
 <212> DNA
 <213> Catalpa sp.

<400> 3
 ttatctttca ggagagattt ttgtttgaat gctcccccg ttgagctttt tgtggaaaaat 60
 taccctccat cttccgtgaa ttgagacccc tgtccatatg gctcaaactg tccgatatgg 120
 gatcctaccc aaatacgact acggcaatcc cagcttcaac ttggcccatt atgggtgaatc 180
 cagacctccc gtttacgatt tatccaagat tcccctcgac attccgctct tcctaagcta 240
 tggaggacaa gatgcattgt cggatgttaa ggatgtcgag acattgctcg atagtctcaa 300
 gttacacgat gtggataagc tgcattgtgca gtatatcaag gattatgctc atgccgactt 360
 cattatcgga gttactgcaa aagatatagt ttataatcag attgtaactt ttttcagaaa 420
 ccaggcttga gaggttcttg attttgaggt gcttttgctg tgagaatgca acagcttgtt 480
 ccactcttgt tgaatgtgaa taagccattt ccgagagatt taatggctgg taaagcttat 540
 tagtttactc atagatacat gtaagaagca acccgataca tagtttgaat cctttatctc 600
 gaaaaggtat tgcattctct cttctacgtc aaaaaaaaaa aaaaata 647

<210> 4
 <211> 116
 <212> PRT
 <213> Catalpa sp.

<400> 4
 Ile Glu Thr Pro Val His Met Ala Gln Thr Val Arg Tyr Gly Ile Leu
 1 5 10 15
 Pro Lys Tyr Asp Tyr Gly Asn Pro Ser Phe Asn Leu Ala His Tyr Gly
 20 25 30
 Glu Ser Arg Pro Pro Val Tyr Asp Leu Ser Lys Ile Pro Leu Asp Ile
 35 40 45
 Pro Leu Phe Leu Ser Tyr Gly Gly Gln Asp Ala Leu Ser Asp Val Lys
 50 55 60
 Asp Val Glu Thr Leu Leu Asp Ser Leu Lys Leu His Asp Val Asp Lys
 65 70 75 80
 Leu His Val Gln Tyr Ile Lys Asp Tyr Ala His Ala Asp Phe Ile Ile
 85 90 95
 Gly Val Thr Ala Lys Asp Ile Val Tyr Asn Gln Ile Val Thr Phe Phe
 100 105 110
 Arg Asn Gln Ala
 115

<210> 5
 <211> 705

<212> DNA
 <213> Catalpa sp.

<220>
 <221> unsure
 <222> (526)

<220>
 <221> unsure
 <222> (561)

<220>
 <221> unsure
 <222> (585)

<220>
 <221> unsure
 <222> (593)

<220>
 <221> unsure
 <222> (664)

<220>
 <221> unsure
 <222> (679)

<400> 5
 gcacgagcca acagcttcct aaatttagct cttctaattc ttctctcatt atcactactc 60
 ctacctcatc aatcattcgc ctccagccgc cgccgttttc ttccgcagaa tgatgtcggt 120
 cttccgcccgc acggcgtttg ctccaccgcc gtaactgtac acggttataa atgccaagaa 180
 tttgaagtaa cgactgatga tggctatata ttaagcgtgc agaggattct ggagggcccg 240
 gccggaggag gagggccgaa gcggccgcgc gttctgctgc aacatggggt tcttgtggac 300
 gggatgacgt ggctggtgaa tggaccggaa caatctttgg cgatgatatt ggctgacaat 360
 ggggttcgacg tctggatttc taacataaga ggaactaggt ttagtcgctg tcatgtcagc 420
 cttgaccta ccgaccta atattgggat tgggcatggg acgatcttgg tgaccacaga 480
 cttaccatcc ctgatcgagt tagtggtcag acaaacgggt cagaanacac actacatagg 540
 gcaatccatg gggaacttta ntagctttgg gatcactttt agganggaaa cangttggca 600
 gggtaaatcg gctgtatggt aagccaattg gctaacgagt catatgcaac tgctctcgag 660
 ttgntagca gatccttgnt ggggaacaca cgatcttggc ctgcg 705

<210> 6
 <211> 157
 <212> PRT
 <213> Catalpa sp.

<400> 6
 Ala Arg Ala Asn Ser Phe Leu Asn Leu Ala Leu Leu Ile Leu Leu Ser
 1 5 10 15
 Leu Ser Leu Leu Leu Pro His Gln Ser Phe Ala Ser Ser Arg Arg Arg
 20 25 30
 Phe Leu Pro Gln Asn Asp Val Val Leu Pro Pro Asp Gly Val Cys Ser
 35 40 45
 Thr Ala Val Thr Val His Gly Tyr Lys Cys Gln Glu Phe Glu Val Thr
 50 55 60
 Thr Asp Asp Gly Tyr Ile Leu Ser Val Gln Arg Ile Leu Glu Gly Arg
 65 70 75 80

Ala Pro Ala Pro Pro Pro Arg Val Pro Leu Pro Pro Ser Arg Trp Ser
20 25 30

Pro Arg Ile Pro Ala His Arg Arg Ala Thr Pro Arg Leu Pro Ala Arg
 35 40 45
 Gly Gly Arg Trp Pro Leu Pro Ala Ala Ala Pro Ala Ala Gly Tyr Pro
 50 55 60
 Cys Thr Glu His Thr Val Gln Thr Asp Asp Gly Phe Leu Leu Ser Leu
 65 70 75 80
 Gln His Ile Pro His Gly Arg Asn Gly Ile Ala Asp Asn Thr Gly Pro
 85 90 95
 Pro Val Phe Leu Gln His Gly Leu Phe Gln Gly Gly Asp Thr Trp Phe
 100 105 110
 Ile Asn Ser Asn Glu Gln Ser Leu Gly Tyr Ile Leu Ala Asp Asn Gly
 115 120 125
 Phe Asp Val Trp Val Gly Asn Val Arg Gly Thr Arg Trp Ser Lys Gly
 130 135 140
 His Ser Thr Leu Ser Val His Asp Lys Leu Phe Trp Asp Trp Ser Trp
 145 150 155 160
 Gln Asp Leu Ala Glu Tyr Asp Val Leu Ala Met Leu Ser Tyr Val Tyr
 165 170 175
 Thr Val Ala Gln Ser Lys Ile Leu Tyr Val Gly His Ser Gln Gly Thr
 180 185 190
 Ile Met Gly Leu Ala Ala Phe Thr Met Pro Glu Thr Val Lys Met Ile
 195 200 205
 Ser Ser Ala Ala Leu Leu Cys Pro Ile Ser Tyr Leu Asp His Val Ser
 210 215 220
 Ala Ser Phe Val Leu Arg Ala Val Ala Met His Leu Asp Glu Met Leu
 225 230 235 240
 Val Ile Met Gly Ile His Gln Leu Asn Phe Arg Ser Asp Met Gly Val
 245 250 255
 Gln Ile Leu Asp Ser Leu Cys Asp Asp Glu His Leu Asp Cys Asn Asp
 260 265 270
 Leu Leu Ser Ser Ile Thr Val Lys Thr Val Val Gln Ser Ser
 275 280 285

<210> 9
 <211> 509
 <212> DNA
 <213> Zea mays

<220>
 <221> unsure
 <222> (162)

<220>
 <221> unsure
 <222> (277)

<220>
 <221> unsure
 <222> (284)

<220>
 <221> unsure
 <222> (290)

<220>
 <221> unsure
 <222> (295)

<220>
 <221> unsure
 <222> (386)

<220>
 <221> unsure
 <222> (406)

<220>
 <221> unsure
 <222> (413)

<220>
 <221> unsure
 <222> (443)

<220>
 <221> unsure
 <222> (468)

<220>
 <221> unsure
 <222> (484)

<220>
 <221> unsure
 <222> (489)

<400> 9
 cgatcgagat ggctcagaag gatctctatc taccgttcct ggctctttcc atcattgcct 60
 gctgcttgat gaacctgcaa agtggttctca gctcaagcag gatgcggaat actacaaacg 120
 atattagtga tgacaaatgc cccccacaaac ctcacccctt angtatgtgc aggtcccagag 180
 tagcagctta cggctatcca tgtgaggaat accatgtgac aacggaggat ggctacatcc 240
 ttagcttaaa gaagatcccc tatggtctct ctggtgncac cganattacn agganaccgg 300
 tactactggt ccatgggcta ctggtggatg gtttctgttg ggtactgaac acacccaaaac 360
 aatcactggg cttcacctgg ctgaangtgg tttgaaattt ggatcnccac tcncccgaaa 420
 aaaatccacc gagggacaca ccnctcccc aaaaaccggc tttgggangg aatggaacac 480
 tgcnaaaana actcccgcgt gctgaatcc 509

<210> 10
 <211> 125
 <212> PRT
 <213> Zea mays

<220>
 <221> UNSURE
 <222> (52)

<220>

<221> UNSURE

<222> (90)

<220>

<221> UNSURE

<222> (92)

<220>

<221> UNSURE

<222> (96)

<400> 10

Met Ala Gln Lys Asp Leu Tyr Leu Pro Phe Leu Ala Leu Ser Ile Ile
 1 5 10 15

Ala Cys Cys Leu Met Asn Leu Gln Ser Val Leu Ser Ser Ser Arg Met
 20 25 30

Arg Asn Thr Thr Asn Asp Ile Ser Asp Asp Lys Cys Pro Pro Gln Pro
 35 40 45

His Pro Leu Xaa Met Cys Arg Ser Arg Val Ala Ala Tyr Gly Tyr Pro
 50 55 60

Cys Glu Glu Tyr His Val Thr Thr Glu Asp Gly Tyr Ile Leu Ser Leu
 65 70 75 80

Lys Lys Ile Pro Tyr Gly Leu Ser Gly Xaa Thr Xaa Ile Thr Arg Xaa
 85 90 95

Pro Val Leu Leu Phe His Gly Leu Leu Val Asp Gly Phe Cys Trp Val
 100 105 110

Leu Asn Thr Pro Lys Gln Ser Leu Gly Phe Thr Trp Leu
 115 120 125

<210> 11

<211> 273

<212> DNA

<213> Zea mays

<220>

<221> unsure

<222> (8)

<220>

<221> unsure

<222> (20)

<220>

<221> unsure

<222> (229)

<220>

<221> unsure

<222> (236)

<220>

<221> unsure

<222> (241)

<220>
 <221> unsure
 <222> (249)

<220>
 <221> unsure
 <222> (268)

<400> 11
 cttcctcntg cacgcttcgn ttccagctct actggaactg gtcctgggat gacctggtag 60
 tcaacgacct gccggccatg gtcgacttcg tcgtcaaaca gaccggccag aagcctcact 120
 acgtcggaca ctccatgggg acgctggtgg cgctggcggc cttctcggag ggccgggtgg 180
 tgagccagct gaaatccgcg gcgctgctca cgccggtggc ctacctcgnc cacatnaaca 240
 nccccaatng gaatcctggt tggccaangc gtt 273

<210> 12
 <211> 90
 <212> PRT
 <213> Zea mays

<220>
 <221> UNSURE
 <222> (76)

<220>
 <221> UNSURE
 <222> (78)

<220>
 <221> UNSURE
 <222> (80)

<220>
 <221> UNSURE
 <222> (83)

<220>
 <221> UNSURE
 <222> (89)

<400> 12
 Ser Ser Cys Thr Leu Arg Phe Gln Leu Tyr Trp Asn Trp Ser Trp Asp
 1 5 10 15
 Asp Leu Val Val Asn Asp Leu Pro Ala Met Val Asp Phe Val Val Lys
 20 25 30
 Gln Thr Gly Gln Lys Pro His Tyr Val Gly His Ser Met Gly Thr Leu
 35 40 45
 Val Ala Leu Ala Ala Phe Ser Glu Gly Arg Val Val Ser Gln Leu Lys
 50 55 60
 Ser Ala Ala Leu Leu Thr Pro Val Ala Tyr Leu Xaa His Xaa Asn Xaa
 65 70 75 80
 Pro Asn Xaa Asn Pro Gly Trp Pro Xaa Arg
 85 90

<210> 13
 <211> 1483
 <212> DNA
 <213> *Oryza sativa*

<400> 13
 gcacgagtag acagcgcggc gggcggtggc gatggcgatg gggggccacg cccccggagg 60
 agcgctcccc ctgacctccc tcgtcgtctc ttgctgcggt cgcacgctct ccggagcctc 120
 ccagcgcgcc gccgccctcc gccgcgtcgg ctccggctcc ggcggcctct gcgaccagct 180
 gctcctgcca ctcggtacc cctgcaccga gcacaacgtt gaaacaaaag atggattcct 240
 tttatctctt cagcatatcc cacatggcaa aaataaagca gcagatagta ctggccctcc 300
 agtttttctt caacatgggtc tttttcaggg aggagacaca tggttcataa actctgctga 360
 gcaatcactt gggatatatcc ttgctgataa cggttttgat gtttggattg ggaatgtccg 420
 tggagcgctg caagagttag ctgaatatga ccttttagca atgctaggct atgtgtatac 480
 agtcacacag tccaaaattc tatatgtggg gcattcacag ggaactataa tgggtttggc 540
 ggctttgacg atgcccgaat tagtaaaaat gattagctct gcagcacttc tttgtcctat 600
 ttcttatctt gatcatgtta gtgctagttt tgttctcaga gcagtcgcca tgcactctga 660
 tcagatgctt gttactatgg gaattcacca gctgaacttc cgtagcgaca tgggggttca 720
 aatagtagat tctttgtgct atggtgaaca cgtggattgc aacaatttgc tatctgcgat 780
 tacaggggaa aactgttgct tcaatacatc aaggattgat tattatttgg agtatgaacc 840
 tcatccatca tcgacaaaaa atctgcacca tcttttccag atgatcagga aaggcacttt 900
 cgcaagtat gactatgggt tattgggaaa cctaaggcgc tacggtcatt tgcgtcctcc 960
 cgcatattgac ctaagcagca taccagaatc actgccata tggatgggat atggaggtct 1020
 tgatgcattg gctgatgtaa ccgatgttca gcgtactatc agagagctgg gatctacacc 1080
 agaacttctg tacattgggt actatggcca tattgatatt gttatgagcg tgaaggcgaa 1140
 agatgatgtt tatgtggacc taataagatt tcttagggaa aatggatggc ataatagcta 1200
 ttaggatgtc ttcattgtga taataaaaac atctgtacag tattggtcct ctcccgatgt 1260
 gagtatgtat atattgcata tgagcttgtt ggatctatgg tgcattgctc caagtctaaa 1320
 acgctgtcag cagcaattgt atcattgtat ccaacttacc gctccactac tgtatatcca 1380
 ttatagaaaa ccctcttcat ttcctcttca aaaaaaaaaa aaa 1440
 1483

<210> 14
 <211> 410
 <212> PRT
 <213> *Oryza sativa*

<400> 14
 Met Ala Met Ala Gly His Ala Pro Gly Gly Ala Leu Pro Leu Ile Leu
 1 5 10 15
 Leu Val Val Ser Cys Cys Gly Arg Ile Val Ser Gly Ala Ser Pro Ala
 20 25 30
 Ala Ala Ala Leu Arg Arg Val Gly Ser Gly Ser Gly Gly Leu Cys Asp
 35 40 45
 Gln Leu Leu Leu Pro Leu Gly Tyr Pro Cys Thr Glu His Asn Val Glu
 50 55 60
 Thr Lys Asp Gly Phe Leu Leu Ser Leu Gln His Ile Pro His Gly Lys
 65 70 75 80
 Asn Lys Ala Ala Asp Ser Thr Gly Pro Pro Val Phe Leu Gln His Gly
 85 90 95
 Leu Phe Gln Gly Gly Asp Thr Trp Phe Ile Asn Ser Ala Glu Gln Ser
 100 105 110
 Leu Gly Tyr Ile Leu Ala Asp Asn Gly Phe Asp Val Trp Ile Gly Asn
 115 120 125

Val Arg Gly Thr Arg Trp Ser Lys Gly His Ser Thr Phe Ser Val His
 130 135 140
 Asp Lys Leu Phe Trp Asp Trp Ser Trp Gln Glu Leu Ala Glu Tyr Asp
 145 150 155 160
 Leu Leu Ala Met Leu Gly Tyr Val Tyr Thr Val Thr Gln Ser Lys Ile
 165 170 175
 Leu Tyr Val Gly His Ser Gln Gly Thr Ile Met Gly Leu Ala Ala Leu
 180 185 190
 Thr Met Pro Glu Ile Val Lys Met Ile Ser Ser Ala Ala Leu Leu Cys
 195 200 205
 Pro Ile Ser Tyr Leu Asp His Val Ser Ala Ser Phe Val Leu Arg Ala
 210 215 220
 Val Ala Met His Leu Asp Gln Met Leu Val Thr Met Gly Ile His Gln
 225 230 235 240
 Leu Asn Phe Arg Ser Asp Met Gly Val Gln Ile Val Asp Ser Leu Cys
 245 250 255
 Asp Gly Glu His Val Asp Cys Asn Asn Leu Leu Ser Ala Ile Thr Gly
 260 265 270
 Glu Asn Cys Cys Phe Asn Thr Ser Arg Ile Asp Tyr Tyr Leu Glu Tyr
 275 280 285
 Glu Pro His Pro Ser Ser Thr Lys Asn Leu His His Leu Phe Gln Met
 290 295 300
 Ile Arg Lys Gly Thr Phe Ala Lys Tyr Asp Tyr Gly Leu Leu Gly Asn
 305 310 315 320
 Leu Arg Arg Tyr Gly His Leu Arg Pro Pro Ala Phe Asp Leu Ser Ser
 325 330 335
 Ile Pro Glu Ser Leu Pro Ile Trp Met Gly Tyr Gly Gly Leu Asp Ala
 340 345 350
 Leu Ala Asp Val Thr Asp Val Gln Arg Thr Ile Arg Glu Leu Gly Ser
 355 360 365
 Thr Pro Glu Leu Leu Tyr Ile Gly Asp Tyr Gly His Ile Asp Phe Val
 370 375 380
 Met Ser Val Lys Ala Lys Asp Asp Val Tyr Val Asp Leu Ile Arg Phe
 385 390 395 400
 Leu Arg Glu Asn Gly Trp His Asn Ser Tyr
 405 410

<210> 15
 <211> 395
 <212> DNA
 <213> Oryza sativa

<220>
<221> unsure
<222> (12)

<220>
<221> unsure
<222> (24)

<220>
<221> unsure
<222> (29)

<220>
<221> unsure
<222> (33)

<220>
<221> unsure
<222> (43)

<220>
<221> unsure
<222> (78)

<220>
<221> unsure
<222> (182)

<220>
<221> unsure
<222> (265)

<220>
<221> unsure
<222> (300)

<220>
<221> unsure
<222> (302)

<220>
<221> unsure
<222> (306)

<220>
<221> unsure
<222> (347)

<220>
<221> unsure
<222> (351)

<220>
<221> unsure
<222> (367)

<220>
<221> unsure
<222> (370)

<220>

<221> unsure

<222> (380)

<220>

<221> unsure

<222> (386)

<400> 15

acatctttca cnggcaaaaa ctantgtcng aanaattcag canccgacat cttcctcaag 60
 tacgagcccc agccaacntc cacaaaaaacc ttgatccatc tcgctcaaac ggtgagagac 120
 ggggtttctga ccaagtacga ctacgtgatg ccggacgcga acgtggccag gtacgggcag 180
 gncgacccgc cggcgtagca catggcggcg atcccggcgt ggttcccat cttcctcagc 240
 tacggcgccc gggactcgct gtccnacccc gccgatcgct gccctcctcc tcgacgaten 300
 cngccncggc ggccacgtcg ggcaccggct catccgtgcc agtaacnttc nccatactcg 360
 cccacgncn acttcgtcan tcgggntttc tgcgc 395

<210> 16

<211> 80

<212> PRT

<213> Oryza sativa

<220>

<221> UNSURE

<222> (8)

<220>

<221> UNSURE

<222> (10)..(11)

<220>

<221> UNSURE

<222> (15)

<220>

<221> UNSURE

<222> (61)

<400> 16

Thr Ser Phe Thr Gly Lys Asn Xaa Cys Xaa Xaa Asn Ser Ala Xaa Asp
 1 5 10 15

Ile Phe Leu Lys Tyr Glu Pro Gln Pro Thr Ser Thr Lys Thr Leu Ile
 20 25 30

His Leu Ala Gln Thr Val Arg Asp Gly Val Leu Thr Lys Tyr Asp Tyr
 35 40 45

Val Met Pro Asp Ala Asn Val Ala Arg Tyr Gly Gln Xaa Asp Pro Pro
 50 55 60

Ala Tyr Asp Met Ala Ala Ile Pro Ala Trp Phe Pro Ile Phe Leu Ser
 65 70 75 80

<210> 17

<211> 1718

<212> DNA

<213> Glycine max

<400> 17

```

ggaatcaaatt attcaactcg ttttcccatc cttttgtgtc tctctttttc cgtttcatac 60
actttttctt taccttttatt gttccaatct tatectatcc tttaaatata cacacacaaa 120
aatacattaa cacttcaatc ccacgctttc aatagataga tagagcattc attcatcacc 180
aacatggctc ttctaggctt aatgagtttt gctgccttga cccttttctt ggtcctaaca 240
actgtgcttc gtcaagcaca cgcttcaagc cgtggcaact taggcagaaa catcaaccct 300
tcagtgtatg gcataatgtgc ctcttctgtc atttgtcatg gatacaagtg tcaagaacac 360
gaggttacaa ctgatgatgg ttacattctg agcctgcaaa ggatcccaga aggtcgaggt 420
aaaaagcagt ggagtgggac aaggaagcaa ccagtgggta tacaacatgg agttcttgta 480
gatggatga catggcttct aaaccaccca gagcaagatc tgccgttgat tttagctgat 540
aatggatttg acgtgtggat tgcaaacaca agaggaacca gatatagtcg ccgacacatc 600
tcattggacc cctctagcca ggcctattgg aattggtctt gggatgaact tgtctcctat 660
gatttccctg cggtgtttta ttatgtgttc agccaaacgg ggcagaagat caattacgtt 720
ggcattcat tgggaacttt ggtagctttg gcacctctct cggaaggaaa attggttacc 780
cagctgaaat cagcagcctt gttgagccct atagcctatt taagccacat gaatacagca 840
cttgggtgtg ttgcacccaa gtcctttgtt ggtgagatca ctaccctctt cggctcagca 900
gaattttaat caaaaagggtt agctgttgat gcctttctca agtctctctg tgctcaccct 960
gggatagact gctatgactt gttgactgca ctaactggta aaaattgctg cctcaattct 1020
tcaactgttg atctattctt gatgaatgag cctcagtcac catcaacaaa gaacatggtg 1080
cacttgctc agactgttag acttggggcg ttgacaaaat tcaattatgt gagaccagac 1140
tataacatta tgactatgg agaaatattt cctccaatct ataacccttc caacatcccc 1200
cacgatctcc ctctcttcat tagctatggg ggaagagatg cactttcaga tgtccgtgat 1260
gttgagaatt tgcttgataa actcaagttc catgatgaga acaagcgag cgttcagttc 1320
atccaggaat atgctcatgc tgactacatt atggggttca atgccaagga cttggtgtat 1380
aatgctgttc tttcattttt caatcatcaa gtttaacact ggatagaatg aatcaagttg 1440
tatgaaaaga gtgccttcat gtattaggta gctatcattg agatcaatct aagtatcta 1500
gtggagatta agtaacggct aattacaaaa gtaatgaagt attatcacta gtgatttgc 1560
ttggtgttgc aaatggctat tgcactatc tattgtgttg cattgtaatg cagaggaaa 1620
tggttttgg cttcagttat ctaagatgaa aaacgtggat gagatcattt atcaaaagaa 1680
ttataaaaac tatgtttcca aaaaaaaaaa aaaaaaaa 1718

```

<210> 18

<211> 410

<212> PRT

<213> Glycine max

<400> 18

```

Met Ala Leu Leu Gly Leu Met Ser Phe Ala Ala Leu Thr Leu Phe Leu
 1             5             10             15

Val Leu Thr Thr Val Pro Arg Gln Ala His Ala Ser Ser Arg Gly Asn
      20             25             30

Leu Gly Arg Asn Ile Asn Pro Ser Val Tyr Gly Ile Cys Ala Ser Ser
      35             40             45

Val Ile Val His Gly Tyr Lys Cys Gln Glu His Glu Val Thr Thr Asp
      50             55             60

Asp Gly Tyr Ile Leu Ser Leu Gln Arg Ile Pro Glu Gly Arg Gly Lys
      65             70             75             80

Ser Ser Gly Ser Gly Thr Arg Lys Gln Pro Val Val Ile Gln His Gly
      85             90             95

Val Leu Val Asp Gly Met Thr Trp Leu Leu Asn Pro Pro Glu Gln Asp
      100            105            110

Leu Pro Leu Ile Leu Ala Asp Asn Gly Phe Asp Val Trp Ile Ala Asn
      115            120            125

```

Thr Arg Gly Thr Arg Tyr Ser Arg Arg His Ile Ser Leu Asp Pro Ser
 130 135 140
 Ser Gln Ala Tyr Trp Asn Trp Ser Trp Asp Glu Leu Val Ser Tyr Asp
 145 150 155 160
 Phe Pro Ala Val Phe Asn Tyr Val Phe Ser Gln Thr Gly Gln Lys Ile
 165 170 175
 Asn Tyr Val Gly His Ser Leu Gly Thr Leu Val Ala Leu Ala Ser Phe
 180 185 190
 Ser Glu Gly Lys Leu Val Thr Gln Leu Lys Ser Ala Ala Leu Leu Ser
 195 200 205
 Pro Ile Ala Tyr Leu Ser His Met Asn Thr Ala Leu Gly Val Val Ala
 210 215 220
 Pro Lys Ser Phe Val Gly Glu Ile Thr Thr Leu Phe Gly Leu Ala Glu
 225 230 235 240
 Phe Asn Pro Lys Gly Leu Ala Val Asp Ala Phe Leu Lys Ser Leu Cys
 245 250 255
 Ala His Pro Gly Ile Asp Cys Tyr Asp Leu Leu Thr Ala Leu Thr Gly
 260 265 270
 Lys Asn Cys Cys Leu Asn Ser Ser Thr Val Asp Leu Phe Leu Met Asn
 275 280 285
 Glu Pro Gln Ser Thr Ser Thr Lys Asn Met Val His Leu Ala Gln Thr
 290 295 300
 Val Arg Leu Gly Ala Leu Thr Lys Phe Asn Tyr Val Arg Pro Asp Tyr
 305 310 315 320
 Asn Ile Met His Tyr Gly Glu Ile Phe Pro Pro Ile Tyr Asn Leu Ser
 325 330 335
 Asn Ile Pro His Asp Leu Pro Leu Phe Ile Ser Tyr Gly Gly Arg Asp
 340 345 350
 Ala Leu Ser Asp Val Arg Asp Val Glu Asn Leu Leu Asp Lys Leu Lys
 355 360 365
 Phe His Asp Glu Asn Lys Arg Ser Val Gln Phe Ile Gln Glu Tyr Ala
 370 375 380
 His Ala Asp Tyr Ile Met Gly Phe Asn Ala Lys Asp Leu Val Tyr Asn
 385 390 395 400
 Ala Val Leu Ser Phe Phe Asn His Gln Val
 405 410

<210> 19

<211> 1438

<212> DNA

<213> Glycine max

<400> 19

```

gcaattcaga ataacaataa aggggtggatg aggatccaga ggttcttggc cacactggcc 60
ataactgtct ccatactctt gggaaatgga aaccccggtc agtgcttcga cggcggtagc 120
caccaaaaac agcaacacag tttgtgtgaa gagctcatta tccctacgg ttaccctgc 180
tccgagcata cgattcaaac gaaggatggg ttcttgtag gtcttcaacg tgtctcttct 240
tcttcttctc ttcggcttcg gaaccatgga gatggaggcc ctccggttct gcttctgcat 300
ggattattca tggcagtgga tgcattggtt ctaaatactc cggaacaatc acttggttc 360
atacttgcaq atcatggtt tgatgtttgg gtaggaaacg tgcgtggaac acgctggagc 420
catggacata tatctttatt agagaagaaa aagcaatttt gggattggag ttggcaggaa 480
ttagccctgt atgatgttc ggaaatgatc aattacatta attcagtaac aaactcaaag 540
atattttag ttgggcattc acaggggaca attatatctt tggctgcctt cactcaacca 600
gagatagtag aaaaggttga ggctgcagct cttctatctc caatatcata cttggatcat 660
gtcagtgcac ctcttgact tagaatggtt aagatgcaca ttgatgagat gattcttacc 720
atgggcattc atcaactaaa cttcaaaagc gaatgggggg ccagtctctt ggtttctcta 780
tgtgataccc gcctaagttg caatgacatg cttcatcca taacagggaa gaattgttgc 840
ttcaatgagt cacgtgtgga gttttatctt gaacaagaac ctcatccatc atcgtctaaa 900
aacttgaacc accttttcca gatgatccgc aaaggtacct actccaagta tgatttga 960
aagctaaaaa atctgataga gtacggcaag ttcaatccac caaagttcga tcttagtcgc 1020
atacccaaat cattgcctct gtggatggct tacgggtgaa atgatgctct ggcagatata 1080
actgatttcc agcacacact caaggaattg ccattccccac cggaagtggg ttatcttgaa 1140
aactatggtc atgttgactt cattttaagc ttgcaagcaa aacaagatct ttatgacct 1200
atgattagtt ttttcaagtc atccggaaaa ttagtagta tgtaatgttt gttctcttcc 1260
ggatgatgg atgtaattac tgtaatggc tacgggtcca tctattactg cacttactgt 1320
aaagttgaaa tattaatatt ctgtggagtc caccttgatt ttctgtatgt atatatgatg 1380
acagatatat aaagatcggc gtcgcatgac ctgtttctgc aaaaaaaaaa aaaaaaaa 1438

```

<210> 20

<211> 405

<212> PRT

<213> Glycine max

<400> 20

```

Met Arg Ile Gln Arg Phe Leu Ala Thr Leu Ala Ile Thr Val Ser Ile
  1             5             10             15

Leu Leu Gly Asn Gly Asn Pro Val Gln Cys Phe Asp Gly Gly Ser His
          20             25             30

Gln Lys Gln Gln His Ser Leu Cys Glu Glu Leu Ile Ile Pro Tyr Gly
          35             40             45

Tyr Pro Cys Ser Glu His Thr Ile Gln Thr Lys Asp Gly Phe Leu Leu
          50             55             60

Gly Leu Gln Arg Val Ser Ser Ser Ser Ser Leu Arg Leu Arg Asn His
          65             70             75             80

Gly Asp Gly Gly Pro Pro Val Leu Leu Leu His Gly Leu Phe Met Ala
          85             90             95

Gly Asp Ala Trp Phe Leu Asn Thr Pro Glu Gln Ser Leu Gly Phe Ile
          100            105            110

Leu Ala Asp His Gly Phe Asp Val Trp Val Gly Asn Val Arg Gly Thr
          115            120            125

Arg Trp Ser His Gly His Ile Ser Leu Leu Glu Lys Lys Lys Gln Phe
          130            135            140

Trp Asp Trp Ser Trp Gln Glu Leu Ala Leu Tyr Asp Val Ala Glu Met
          145            150            155            160

```

Ile Asn Tyr Ile Asn Ser Val Thr Asn Ser Lys Ile Phe Val Val Gly
 165 170 175
 His Ser Gln Gly Thr Ile Ile Ser Leu Ala Ala Phe Thr Gln Pro Glu
 180 185 190
 Ile Val Glu Lys Val Glu Ala Ala Ala Leu Leu Ser Pro Ile Ser Tyr
 195 200 205
 Leu Asp His Val Ser Ala Pro Leu Val Leu Arg Met Val Lys Met His
 210 215 220
 Ile Asp Glu Met Ile Leu Thr Met Gly Ile His Gln Leu Asn Phe Lys
 225 230 235 240
 Ser Glu Trp Gly Ala Ser Leu Leu Val Ser Leu Cys Asp Thr Arg Leu
 245 250 255
 Ser Cys Asn Asp Met Leu Ser Ser Ile Thr Gly Lys Asn Cys Cys Phe
 260 265 270
 Asn Glu Ser Arg Val Glu Phe Tyr Leu Glu Gln Glu Pro His Pro Ser
 275 280 285
 Ser Ser Lys Asn Leu Asn His Leu Phe Gln Met Ile Arg Lys Gly Thr
 290 295 300
 Tyr Ser Lys Tyr Asp Tyr Gly Lys Leu Lys Asn Leu Ile Glu Tyr Gly
 305 310 315 320
 Lys Phe Asn Pro Pro Lys Phe Asp Leu Ser Arg Ile Pro Lys Ser Leu
 325 330 335
 Pro Leu Trp Met Ala Tyr Gly Gly Asn Asp Ala Leu Ala Asp Ile Thr
 340 345 350
 Asp Phe Gln His Thr Leu Lys Glu Leu Pro Ser Pro Pro Glu Val Val
 355 360 365
 Tyr Leu Glu Asn Tyr Gly His Val Asp Phe Ile Leu Ser Leu Gln Ala
 370 375 380
 Lys Gln Asp Leu Tyr Asp Pro Met Ile Ser Phe Phe Lys Ser Ser Gly
 385 390 395 400
 Lys Phe Ser Ser Met
 405

<210> 21

<211> 737

<212> DNA

<213> Zea mays

<400> 21

gcacgagggt ttgtgccctt gatctatctg ttaaatttgg atcgcaggag gttgaactca 60
 tgaactttgg acagcctcgg ataggcaatc ctgcatttgc tgtatacttt ggtgaacaag 120
 tccaagaac aatccgtgtg acccatcaga atgatattgt gccgcattta ccaccgtatt 180
 attattacct aggtgaatgg acataccacc acttcgctag agagggttgg cttcatgaga 240
 gcatagatgg aaatgtagtt accagaaacg agacggtatg tgatgattct ggtgaagacc 300
 cgacctgtag caggctcggtc tatgggatga gcgtagcaga tcatcttgag tactatgatg 360

```

tcacactaca tgctgattca agaggaacct gtcaattcgt gattggtgca gccaaccaag 420
tatacaacta cgttcgtgaa gttgatggat ccatcatcct gtcaagatac ccgcaagaac 480
cacaagctct agaatctatg tgactttgta tgccacggaa tgcacccctg tacagtattt 540
ttcattttca ttttgtgtac agctcatgaa atgctgggcg ctctggagc tctccagagg 600
ataaggagag gctcaccctt ttaaatgtgc cccctttgct caagtgaagaa tcgtgcatgt 660
aagctccata agattgtccg cacaattcaa tttgtgtata taaataatac tatgtgttac 720
taaaaaaaaa aaaaaaa 737

```

<210> 22
 <211> 166
 <212> PRT
 <213> Zea mays

<400> 22
 Thr Arg Phe Cys Ala Leu Asp Leu Ser Val Lys Phe Gly Ser Gln Glu
 1 5 10 15
 Val Glu Leu Met Thr Phe Gly Gln Pro Arg Ile Gly Asn Pro Ala Phe
 20 25 30
 Ala Val Tyr Phe Gly Glu Gln Val Pro Arg Thr Ile Arg Val Thr His
 35 40 45
 Gln Asn Asp Ile Val Pro His Leu Pro Pro Tyr Tyr Tyr Tyr Leu Gly
 50 55 60
 Glu Trp Thr Tyr His His Phe Ala Arg Glu Val Trp Leu His Glu Ser
 65 70 75 80
 Ile Asp Gly Asn Val Val Thr Arg Asn Glu Thr Val Cys Asp Asp Ser
 85 90 95
 Gly Glu Asp Pro Thr Cys Ser Arg Ser Val Tyr Gly Met Ser Val Ala
 100 105 110
 Asp His Leu Glu Tyr Tyr Asp Val Thr Leu His Ala Asp Ser Arg Gly
 115 120 125
 Thr Cys Gln Phe Val Ile Gly Ala Ala Asn Gln Val Tyr Asn Tyr Val
 130 135 140
 Arg Glu Val Asp Gly Ser Ile Ile Leu Ser Arg Tyr Pro Gln Glu Pro
 145 150 155 160
 Gln Ala Leu Glu Ser Met
 165

<210> 23
 <211> 1434
 <212> DNA
 <213> Zea mays

<220>
 <221> unsure
 <222> (226)

<220>
 <221> unsure
 <222> (315)

<220>
 <221> unsure
 <222> (1306)

<220>
 <221> unsure
 <222> (1349)

<220>
 <221> unsure
 <222> (1359)

<220>
 <221> unsure
 <222> (1368)

<220>
 <221> unsure
 <222> (1373)

<400> 23
 acccacgcgt ccgcccacgc gtccggctct ggaagcaggt tcagatttag cctgggtgcg 60
 tctgcaggtt ccggttcacg gagagatgga gcttgggtgc caaagtggta gctctcgac 120
 tcttgctgtc tgctgcttct catggaagag agttgcctgt caagagtagt gaccgcagtt 180
 ttatctacaa ccatactctt gcaaagacgc ttgtggaata tgcathagcg gtgtatatga 240
 cagatttaac cgctctgttt acgtggacat gctcaagatg caatgacttg actcaaggat 300
 tcgagatgag atccntaatt gttgatgtgg agaaactgct tgcaggcatt gttggtgtag 360
 atcatagtct gaattcgata attgttgcaa tcaggggaac tcaagagaac agtgtacaga 420
 attggataaa agacttgata tggaagcagc ttgatctaag tnatccaaac atgccaatg 480
 caaagtgca cagtggtatt ttctctcgt ataacaatac aattttgcgt ctacttatca 540
 caagtgtgtg gcacaaggca agaaagtcac atggagatat caatgtcata gtgacaggcc 600
 actcgatggg aggagctatg gcttcttttt gcgcgctcga tcttgctatg aagcttggag 660
 gtggcagtg gcaactcatg acttttgggc agcctcgtgt tggcaatgct gcattcgctt 720
 catacttcgc caaatatgta cccaacacaa ttcgagtgac acacgggcat gatattgtgc 780
 cacatttggc accttatttc tctttcttc cccagctgac ataccaccat ttccaagag 840
 aggtatgggt ccaggattct gatggcaaca caactgaacg gatttgtgac gacagcgggtg 900
 aagaccaga ttgttgacag tgcatctcca tgttcggctt gaggattcag gaccattcac 960
 ttacctagga gttgatatgg aagcggacga ctggagcacc tgtagaatca tcacagctca 1020
 aagggttcag cagttccgac tggagctagc ggcaacatca tgatgaccaa gcacgatatc 1080
 gacgtctcca tcgttgaacc tagtgtacaa aacagattgg agcagttcta gataggcgga 1140
 acattcgttt tgtccagatt cagagaagca acagcctctt gttaatgcag tggaaattgt 1200
 tcagagacga aggcgaccct tgtttctcta ttagttcggc cagaatggga tgttctttca 1260
 gtcaaggaat aaatcgggag catctcttgt aacaaagaga tcaganatga tgtcataagg 1320
 aaaatcatag gacgtttatg ctgattggna ggattgctnt ggtaatana gancatgtaa 1380
 cttcatgctt attcagaaca tagaccagct actgaaatta gttacgaaaa 1434

<210> 24
 <211> 296
 <212> PRT
 <213> Zea mays

<220>
 <221> UNSURE
 <222> (50)

<220>
 <221> UNSURE
 <222> (80)

<220>

<221> UNSURE

<222> (129)

<400> 24

Met Glu Arg Trp Ser Leu Gly Ala Lys Val Val Ala Leu Ala Leu Leu
 1 5 10 15
 Leu Ser Ala Ala Ser His Gly Arg Glu Leu Pro Val Lys Ser Ser Asp
 20 25 30
 Arg Ser Phe Ile Tyr Asn His Thr Leu Ala Lys Thr Leu Val Glu Tyr
 35 40 45
 Ala Xaa Ala Val Tyr Met Thr Asp Leu Thr Ala Leu Phe Thr Trp Thr
 50 55 60
 Cys Ser Arg Cys Asn Asp Leu Thr Gln Gly Phe Glu Met Arg Ser Xaa
 65 70 75 80
 Ile Val Asp Val Glu Lys Leu Leu Ala Gly Ile Val Gly Val Asp His
 85 90 95
 Ser Leu Asn Ser Ile Ile Val Ala Ile Arg Gly Thr Gln Glu Asn Ser
 100 105 110
 Val Gln Asn Trp Ile Lys Asp Leu Ile Trp Lys Gln Leu Asp Leu Ser
 115 120 125
 Xaa Pro Asn Met Pro Asn Ala Lys Val His Ser Gly Phe Phe Ser Ser
 130 135 140
 Tyr Asn Asn Thr Ile Leu Arg Leu Ala Ile Thr Ser Ala Val His Lys
 145 150 155 160
 Ala Arg Lys Ser Tyr Gly Asp Ile Asn Val Ile Val Thr Gly His Ser
 165 170 175
 Met Gly Gly Ala Met Ala Ser Phe Cys Ala Leu Asp Leu Ala Met Lys
 180 185 190
 Leu Gly Gly Gly Ser Val Gln Leu Met Thr Phe Gly Gln Pro Arg Val
 195 200 205
 Gly Asn Ala Ala Phe Ala Ser Tyr Phe Ala Lys Tyr Val Pro Asn Thr
 210 215 220
 Ile Arg Val Thr His Gly His Asp Ile Val Pro His Leu Pro Pro Tyr
 225 230 235 240
 Phe Ser Phe Leu Pro Gln Leu Thr Tyr His His Phe Pro Arg Glu Val
 245 250 255
 Trp Val Gln Asp Ser Asp Gly Asn Thr Thr Glu Arg Ile Cys Asp Asp
 260 265 270
 Ser Gly Glu Asp Pro Asp Cys Cys Arg Cys Ile Ser Met Phe Gly Leu
 275 280 285
 Arg Ile Gln Asp His Ser Leu Thr
 290 295

<210> 25
 <211> 1560
 <212> DNA
 <213> Zea mays

<220>
 <221> unsure
 <222> (601)

<400> 25
 tcccactgaa ccgggaggcg cccaacttcc ggtccatgat cgggatgata gacgggagga 60
 cggagctgaa gccgctgcca gccagcggcg gccagagga ccggcggctg caggtggtgg 120
 gcgtctccgc cgccgcggga tccggcgcga tgactactac ttggacgtgg agagcggcag 180
 gcggcgggtgc cgctggtgca gcagcagtac gtgaacgggc ggctcgtccg cctccgcacc 240
 ttctccgtgt tcgaggtgag catgatggcc gccaaagatcg cctacgagaa cgccgcctac 300
 atcgagaacg tcgtcaacaa cgtctggaag ttccacttcg tggggttcta caactgctgg 360
 aacaagttcg tgggcgacca cacgacgcag gcgttcgtgt tcaccgacaa ggcaagagga 420
 cgcgagcgtg gtggtggtgt cgttccgggg cacggagccc ttcaacatgc gggactggtc 480
 cacggaacgt aacctgtcgt ggctgggcat gggcgagctg ggccacgtcc acgtcggctt 540
 cctcaaggcg ctgggcctgc aggaggagga cggcaaggac gccacgcggg cgttcccca 600
 nggcgcccc aacgcgctcc cgggcaagcc gctggcctac tacgcgctgc gcgaggaggt 660
 ccagaagcag ctgcagaagc acccgaacgc caacgtcgtg gtcaccggcc acagcctcgg 720
 cgccgcgctg gcgaccatct tcccggcgct gctggcgctt cacggggagc ggggcgtcct 780
 ggaccgcctg ctctccgtgg tcacctacgg gcagccgcgc gtgggcgaca aggtgttcgc 840
 gggctacgtg cgcgccaacg tgcccgtgga gccgctccgg gtggtgtacc gctacgacgt 900
 ggtcccgcgc gtgcccttcg acgcgcgcgc cgtcgccgac ttcgcgcacg gcggcacctg 960
 cgtctacttc gacggctggt acaagggccg cgagatcgcc aaggcgggcg acgcgcccc 1020
 caagaactac ttcgacccca ggtacctgct gtccatgtac ggcaacgcgt ggggggacct 1080
 cttcaagggc gccttctgt gggccaagga gggcaaggac taccgcgagg gcgccgtctc 1140
 gctgtcttac cgcgccaccg gcctgtcgt gcccgcatc gcgtcgaca gccccaggga 1200
 ctacgtcaac gccgtccgcc tcggcagcgt cgcctcggcg tagcttttgg attgcatgtt 1260
 cgtttccatg catgtgtatc attgcatgca ataattggat gaaataaaca gcaataagct 1320
 tcatcagtat tattattgtt gttgttgaat atatgcatcc tctcctctct atatagaatt 1380
 atagatacat gaggcctggc cggccgcgca cgttgctgaa cagttgaagc gcttcccaaa 1440
 aaaaaatgta tcaactgtga agcatatata tccatgcatg catgtgtgcc cgaaattttt 1500
 gtttttaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaacaaa aaaaaaaaaa 1560

<210> 26
 <211> 258
 <212> PRT
 <213> Zea mays

<220>
 <221> UNSURE
 <222> (45)

<400> 26
 Met Arg Asp Trp Ser Thr Asp Val Asn Leu Ser Trp Leu Gly Met Gly
 1 5 10 15
 Glu Leu Gly His Val His Val Gly Phe Leu Lys Ala Leu Gly Leu Gln
 20 25 30
 Glu Glu Asp Gly Lys Asp Ala Thr Arg Ala Phe Pro Xaa Gly Ala Pro
 35 40 45
 Asn Ala Val Pro Gly Lys Pro Leu Ala Tyr Tyr Ala Leu Arg Glu Glu
 50 55 60

Val Gln Lys Gln Leu Gln Lys His Pro Asn Ala Asn Val Val Val Thr
 65 70 75 80
 Gly His Ser Leu Gly Ala Ala Leu Ala Thr Ile Phe Pro Ala Leu Leu
 85 90 95
 Ala Phe His Gly Glu Arg Gly Val Leu Asp Arg Leu Leu Ser Val Val
 100 105 110
 Thr Tyr Gly Gln Pro Arg Val Gly Asp Lys Val Phe Ala Gly Tyr Val
 115 120 125
 Arg Ala Asn Val Pro Val Glu Pro Leu Arg Val Val Tyr Arg Tyr Asp
 130 135 140
 Val Val Pro Arg Val Pro Phe Asp Ala Pro Pro Val Ala Asp Phe Ala
 145 150 155 160
 His Gly Gly Thr Cys Val Tyr Phe Asp Gly Trp Tyr Lys Gly Arg Glu
 165 170 175
 Ile Ala Lys Gly Gly Asp Ala Pro Asn Lys Asn Tyr Phe Asp Pro Arg
 180 185 190
 Tyr Leu Leu Ser Met Tyr Gly Asn Ala Trp Gly Asp Leu Phe Lys Gly
 195 200 205
 Ala Phe Leu Trp Ala Lys Glu Gly Lys Asp Tyr Arg Glu Gly Ala Val
 210 215 220
 Ser Leu Leu Tyr Arg Ala Thr Gly Leu Leu Val Pro Gly Ile Ala Ser
 225 230 235 240
 His Ser Pro Arg Asp Tyr Val Asn Ala Val Arg Leu Gly Ser Val Ala
 245 250 255

Ser Ala

<210> 27
 <211> 432
 <212> DNA
 <213> *Oryza sativa*

<220>
 <221> unsure
 <222> (7)

<220>
 <221> unsure
 <222> (15)

<220>
 <221> unsure
 <222> (27)

<220>
 <221> unsure
 <222> (38)

<220>
 <221> unsure
 <222> (50)

<220>
 <221> unsure
 <222> (94)

<220>
 <221> unsure
 <222> (99)

<220>
 <221> unsure
 <222> (103)

<220>
 <221> unsure
 <222> (105)

<220>
 <221> unsure
 <222> (117)

<400> 27
 catagtnata atacnaacag ttgcggnat tgagattntt ggaaatctgn tcggtgggca 60
 aggaagacat atggaaggct acctataaat gttntagnt cantncgatg ggagggncct 120
 tttagcatcg ttcttgtgcc cttgacctct cttgttaagt atggatcgca ggaagttcaa 180
 ctcagtactt ttggacagcc tcgggtaggc aatccttctt ttgctgcgta cttcagtgc 240
 caagtcccga gaacaatccg tgtgacccat cagaatgaca ttgtcccaca cttgccacca 300
 tatttttgct accttggcga atggacatat caccacttct cgagagaggt ttggcttcat 360
 gagaccatag taggaaatgt agttactagg aatgagacca tctgtgatgg atcaggcgag 420
 gacccaacat gc 432

<210> 28
 <211> 106
 <212> PRT
 <213> Oryza sativa

<400> 28
 Gly Pro Phe Ser Ile Val Leu Val Pro Leu Thr Ser Leu Val Lys Tyr
 1 5 10 15
 Gly Ser Gln Glu Val Gln Leu Met Thr Phe Gly Gln Pro Arg Val Gly
 20 25 30
 Asn Pro Ser Phe Ala Ala Tyr Phe Ser Asp Gln Val Pro Arg Thr Ile
 35 40 45
 Arg Val Thr His Gln Asn Asp Ile Val Pro His Leu Pro Pro Tyr Phe
 50 55 60
 Cys Tyr Leu Gly Glu Trp Thr Tyr His His Phe Ser Arg Glu Val Trp
 65 70 75 80
 Leu His Glu Thr Ile Val Gly Asn Val Val Thr Arg Asn Glu Thr Ile
 85 90 95
 Cys Asp Gly Ser Gly Glu Asp Pro Thr Cys
 100 105

<210> 29
 <211> 1234
 <212> DNA
 <213> Glycine max

<400> 29
 ccactggaag atggaattcg tgagattttt tgattgctgg gaatgatttt caagaaaagg 60
 ccacaaccga agtcttgatt gttttggaca agcatgagaa ccgcgatact tatgtggtag 120
 ctttccgagg aacggaaccc ttgatgcag atgcatgggt cactgacctt gacatctcat 180
 ggtacgcatt cccggcattg gaaaaatgca tgggtggcttc atgaaagcct tagggctaca 240
 gaaaaatgtg ggggtggccta aggagattca aagggatgaa aatcttcccc cgttggccta 300
 ctatgttatt agggacattc taaggaaaagg tttgagtga aatcctaatag caaagtttat 360
 cattacgggt catagtttgg gtggagcact cgcaatcttg taccctacga tcatgttctt 420
 gcatgatgag aagttgctga ttgagagggt ggaagggatc tacacgtttg ggcaaccaag 480
 agttggagat gaagcatatg cacagtatat gagacaaaaa ttgagggaaa attctatcag 540
 gtattgcagg tttgtttatt gcaatgacat agttccgagg ttgccctatg atgataagga 600
 cttgctcttc aagcactttg ggatctgcct tttctttaac aggcgctatg aactcaggat 660
 tctcgaagaa gagccgaata agaactatct ctcgccatgg tgtgtgatac ccatgatgtt 720
 caatgctgtt ttggaactaa taaggagcct taccatagcg tacaaaaatg gacctcacta 780
 tagagaagga tgggttctct ttagtttcag gttgggttgg ctgctgattc ctggcttacc 840
 tgctcacggt ccacaagatt atattaattc cactcttctg ggatcaattg aaaaacattt 900
 taaagcagat tgatgtgtcc gtatacatga tcattccata ccactacgta catgtgtatg 960
 gtcatgcaga ctaaaattta cataatcaag atttttagtt ttgaaaaaaa tggtataaac 1020
 acttgattat gtatcatgtg aagaatagtt atgtatcata atgatcatga ataataaac 1080
 agtttgcgt cagtacgagt tattgtatag taattaataa gctaggttta aagttgtttc 1140
 ctttgggtgca tggatttatc attaatgaga tcaatgtgaa gtttgtttat ttcaaaaaaa 1200
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 1234

<210> 30
 <211> 246
 <212> PRT
 <213> Glycine max

<400> 30
 His Leu Met Val Arg Ile Pro Gly Ile Gly Lys Met His Gly Gly Phe
 1 5 10 15
 Met Lys Ala Leu Gly Leu Gln Lys Asn Val Gly Trp Pro Lys Glu Ile
 20 25 30
 Gln Arg Asp Glu Asn Leu Pro Pro Leu Ala Tyr Tyr Val Ile Arg Asp
 35 40 45
 Ile Leu Arg Lys Gly Leu Ser Glu Asn Pro Asn Ala Lys Phe Ile Ile
 50 55 60
 Thr Gly His Ser Leu Gly Gly Ala Leu Ala Ile Leu Tyr Pro Thr Ile
 65 70 75 80
 Met Phe Leu His Asp Glu Lys Leu Leu Ile Glu Arg Leu Glu Gly Ile
 85 90 95
 Tyr Thr Phe Gly Gln Pro Arg Val Gly Asp Glu Ala Tyr Ala Gln Tyr
 100 105 110
 Met Arg Gln Lys Leu Arg Glu Asn Ser Ile Arg Tyr Cys Arg Phe Val
 115 120 125
 Tyr Cys Asn Asp Ile Val Pro Arg Leu Pro Tyr Asp Asp Lys Asp Leu
 130 135 140

Leu Phe Lys His Phe Gly Ile Cys Leu Phe Phe Asn Arg Arg Tyr Glu
 145 150 155 160
 Leu Arg Ile Leu Glu Glu Glu Pro Asn Lys Asn Tyr Phe Ser Pro Trp
 165 170 175
 Cys Val Ile Pro Met Met Phe Asn Ala Val Leu Glu Leu Ile Arg Ser
 180 185 190
 Phe Thr Ile Ala Tyr Lys Asn Gly Pro His Tyr Arg Glu Gly Trp Phe
 195 200 205
 Leu Phe Ser Phe Arg Leu Val Gly Leu Leu Ile Pro Gly Leu Pro Ala
 210 215 220
 His Gly Pro Gln Asp Tyr Ile Asn Ser Thr Leu Leu Gly Ser Ile Glu
 225 230 235 240
 Lys His Phe Lys Ala Asp
 245

<210> 31
 <211> 490
 <212> DNA
 <213> Glycine max

<400> 31
 gcacgaggag agatggccta aagaaattga aaccgatgag aaccgtccac gtgtctatta 60
 ctccataagg gatttgctaa agaagtgttt gaatagaaat gataaagcaa agtttattct 120
 tacgggtcat agtcttggtg gagcacttgc aattcttttt cccgctatgc taattttgca 180
 tgctgagaca tttcttttgg aaaggcttga aggggtgtac acatttggac agcctagggt 240
 tggagatgaa acatttgcta aatacatgga aaatcaattg aaacattatg gcattaagta 300
 ttttaggttt gtttactgca acgatattgt tcctagggtt ccctttgatg aagatatcat 360
 gaaatttgag ctttttgga catgtcttta ttatgacagg agctatacat gcaaggtaca 420
 tatataagta ttttaatttt ttgattcatg catatatctg tcattgtaat caactttttt 480
 ttttctgggg 490

<210> 32
 <211> 141
 <212> PRT
 <213> Glycine max

<400> 32
 His Glu Glu Arg Trp Pro Lys Glu Ile Glu Thr Asp Glu Asn Arg Pro
 1 5 10 15
 Arg Val Tyr Tyr Ser Ile Arg Asp Leu Leu Lys Lys Cys Leu Asn Arg
 20 25 30
 Asn Asp Lys Ala Lys Phe Ile Leu Thr Gly His Ser Leu Gly Gly Ala
 35 40 45
 Leu Ala Ile Leu Phe Pro Ala Met Leu Ile Leu His Ala Glu Thr Phe
 50 55 60
 Leu Leu Glu Arg Leu Glu Gly Val Tyr Thr Phe Gly Gln Pro Arg Val
 65 70 75 80
 Gly Asp Glu Thr Phe Ala Lys Tyr Met Glu Asn Gln Leu Lys His Tyr
 85 90 95

Gly Ile Lys Tyr Phe Arg Phe Val Tyr Cys Asn Asp Ile Val Pro Arg
 100 105 110

Leu Pro Phe Asp Glu Asp Ile Met Lys Phe Glu His Phe Gly Thr Cys
 115 120 125

Leu Tyr Tyr Asp Arg Ser Tyr Thr Cys Lys Val His Ile
 130 135 140

<210> 33
 <211> 774
 <212> DNA
 <213> Triticum aestivum

<400> 33
 gcacgagaat attcccatca tggtagacagg acattccatg ggagggggcca tggcttcggt 60
 ttgtgccctt gatcttattg tcaactatgg gttaaaggac gtgaccctgc tgacatttgg 120
 gcaacctcgg attggtaatg ctgtgtttgc taccacttt aagaaatact tgccaaacgc 180
 aattcgagtt accaacgcac atgatattgt gcctcatcta ccccggtact accagtactt 240
 cccacagaat acctaccatc atttcccacc agagggttgg gttcataaca ttggactcga 300
 tagcctacta taccgatcgc agcacatctg tgatcattct ggagaaagac cccacttgca 360
 gcaggccctt ggttggaat agcgtccagg cccatacccc ctttcttggc tccagcatcc 420
 atcccagtc gcgcggatca tccagaatcg tcacggatga caatatgctc aggcacaaag 480
 ttgcccctgt agacggtgtt attgtcttct cgaagcagcc tggtttatca gttggtcagc 540
 tactcagtac acagtaaaca agctcaagat tacatggatt tattttgatg ttttttttg 600
 ccaaaqaaca atattcttgt tggcaatcaa agcactatct catgtatata tacgcgtgtg 660
 atcttggtg gattaaatta tcctagctga ggggtgtatt ctgaaatgta caaacatatt 720
 tatgctgatt aaaaaaaaaa aaaaaaatac ttgaggcggc cccgtaccaa aaat 774

<210> 34
 <211> 126
 <212> PRT
 <213> Triticum aestivum

<400> 34
 His Glu Asn Ile Pro Ile Met Val Thr Gly His Ser Met Gly Gly Ala
 1 5 10 15
 Met Ala Ser Phe Cys Ala Leu Asp Leu Ile Val Asn Tyr Gly Leu Lys
 20 25 30
 Asp Val Thr Leu Leu Thr Phe Gly Gln Pro Arg Ile Gly Asn Ala Val
 35 40 45
 Phe Ala Thr His Phe Lys Lys Tyr Leu Pro Asn Ala Ile Arg Val Thr
 50 55 60
 Asn Ala His Asp Ile Val Pro His Leu Pro Pro Tyr Tyr Gln Tyr Phe
 65 70 75 80
 Pro Gln Asn Thr Tyr His His Phe Pro Pro Glu Val Trp Val His Asn
 85 90 95
 Ile Gly Leu Asp Ser Leu Leu Tyr Pro Ile Glu His Ile Cys Asp His
 100 105 110
 Ser Gly Glu Arg Pro His Leu Gln Gln Ala Leu Gly Trp Lys
 115 120 125

<210> 35
 <211> 398

<212> PRT

<213> Canis familiaris

<400> 35

```

Met Trp Leu Leu Leu Thr Ala Ala Ser Val Ile Ser Thr Leu Gly Thr
  1           5           10           15

Thr His Gly Leu Phe Gly Lys Leu His Pro Thr Asn Pro Glu Val Thr
          20           25           30

Met Asn Ile Ser Gln Met Ile Thr Tyr Trp Gly Tyr Pro Ala Glu Glu
          35           40           45

Tyr Glu Val Val Thr Glu Asp Gly Tyr Ile Leu Gly Ile Asp Arg Ile
  50           55           60

Pro Tyr Gly Arg Lys Asn Ser Glu Asn Ile Gly Arg Arg Pro Val Ala
  65           70           75           80

Phe Leu Gln His Gly Leu Leu Ala Ser Ala Thr Asn Trp Ile Ser Asn
          85           90           95

Leu Pro Asn Asn Ser Leu Ala Phe Ile Leu Ala Asp Ala Gly Tyr Asp
          100          105          110

Val Trp Leu Gly Asn Ser Arg Gly Asn Thr Trp Ala Arg Arg Asn Leu
          115          120          125

Tyr Tyr Ser Pro Asp Ser Val Glu Phe Trp Ala Phe Ser Phe Asp Glu
          130          135          140

Met Ala Lys Tyr Asp Leu Pro Ala Thr Ile Asp Phe Ile Leu Lys Lys
          145          150          155          160

Thr Gly Gln Asp Lys Leu His Tyr Val Gly His Ser Gln Gly Thr Thr
          165          170          175

Ile Gly Phe Ile Ala Phe Ser Thr Asn Pro Lys Leu Ala Lys Arg Ile
          180          185          190

Lys Thr Phe Tyr Ala Leu Ala Pro Val Ala Thr Val Lys Tyr Thr Glu
          195          200          205

Thr Leu Leu Asn Lys Leu Met Leu Val Pro Ser Phe Leu Phe Lys Leu
          210          215          220

Ile Phe Gly Asn Lys Ile Phe Tyr Pro His His Phe Phe Asp Gln Phe
          225          230          235          240

Leu Ala Thr Glu Val Cys Ser Arg Glu Thr Val Asp Leu Leu Cys Ser
          245          250          255

Asn Ala Leu Phe Ile Ile Cys Gly Phe Asp Thr Met Asn Leu Asn Met
          260          265          270

Ser Arg Leu Asp Val Tyr Leu Ser His Asn Pro Ala Gly Thr Ser Val
          275          280          285

Gln Asn Val Leu His Trp Ser Gln Ala Val Lys Ser Gly Lys Phe Gln
          290          295          300

```

Ala Phe Asp Trp Gly Ser Pro Val Gln Asn Met Met His Tyr His Gln
 305 310 315 320

Ser Met Pro Pro Tyr Tyr Asn Leu Thr Asp Met His Val Pro Ile Ala
 325 330 335

Val Trp Asn Gly Gly Asn Asp Leu Leu Ala Asp Pro His Asp Val Asp
 340 345 350

Leu Leu Leu Ser Lys Leu Pro Asn Leu Ile Tyr His Arg Lys Ile Pro
 355 360 365

Pro Tyr Asn His Leu Asp Phe Ile Trp Ala Met Asp Ala Pro Gln Ala
 370 375 380

Val Tyr Asn Glu Ile Val Ser Met Met Gly Thr Asp Asn Lys
 385 390 395

<210> 36
 <211> 403
 <212> PRT
 <213> Caenorhabditis elegans

<400> 36
 Met Trp Arg Phe Ala Val Phe Leu Ala Ala Phe Phe Val Gln Asp Val
 1 5 10 15

Val Gly Ser His Gly Asp Pro Glu Leu His Met Thr Thr Pro Gln Ile
 20 25 30

Ile Glu Arg Trp Gly Tyr Pro Ala Met Ile Tyr Thr Val Ala Thr Asp
 35 40 45

Asp Gly Tyr Ile Leu Glu Met His Arg Ile Pro Phe Gly Lys Thr Asn
 50 55 60

Val Thr Trp Pro Asn Gly Lys Arg Pro Val Val Phe Met Gln His Gly
 65 70 75 80

Leu Leu Cys Ala Ser Ser Asp Trp Val Val Asn Leu Pro Asp Gln Ser
 85 90 95

Ala Gly Phe Leu Phe Ala Asp Ala Gly Phe Asp Val Trp Leu Gly Asn
 100 105 110

Met Arg Gly Asn Thr Tyr Ser Met Lys His Lys Asp Leu Lys Pro Ser
 115 120 125

His Ser Ala Phe Trp Asp Trp Ser Trp Asp Glu Met Ala Thr Tyr Asp
 130 135 140

Leu Asn Ala Met Ile Asn His Val Leu Glu Val Thr Gly Gln Asp Ser
 145 150 155 160

Val Tyr Tyr Met Gly His Ser Gln Gly Thr Leu Thr Met Phe Ser His
 165 170 175

Leu Ser Lys Asp Asp Gly Ser Phe Ala Lys Lys Ile Lys Lys Phe Phe
 180 185 190

Ala Leu Ala Pro Ile Gly Ser Val Lys His Ile Lys Gly Phe Leu Ser
 195 200 205
 Phe Phe Ala Asn Tyr Phe Ser Leu Glu Phe Asp Gly Trp Phe Asp Ile
 210 215 220
 Phe Gly Ala Gly Glu Phe Leu Pro Asn Asn Trp Ala Met Lys Leu Ala
 225 230 235 240
 Ala Lys Asp Ile Cys Gly Gly Leu Lys Val Glu Ala Asp Leu Cys Asp
 245 250 255
 Asn Val Leu Phe Leu Ile Ala Gly Pro Glu Ser Asp Gln Trp Asn Gln
 260 265 270
 Thr Arg Val Pro Val Tyr Ala Thr His Asp Pro Ala Gly Thr Ser Thr
 275 280 285
 Gln Asn Ile Val His Trp Met Gln Met Val His His Gly Gly Val Pro
 290 295 300
 Ala Tyr Asp Trp Gly Thr Lys Thr Asn Lys Lys Lys Tyr Gly Gln Ala
 305 310 315 320
 Asn Pro Pro Glu Tyr Asp Phe Thr Ala Ile Lys Gly Thr Lys Ile Tyr
 325 330 335
 Leu Tyr Trp Ser Asp Ala Asp Trp Leu Ala Asp Thr Pro Asp Val Pro
 340 345 350
 Asp Tyr Leu Leu Thr Arg Leu Asn Pro Ala Ile Val Ala Gln Asn Asn
 355 360 365
 His Leu Pro Asp Tyr Asn His Leu Asp Phe Thr Trp Gly Leu Arg Ala
 370 375 380
 Pro Asp Asp Ile Tyr Arg Pro Ala Ile Lys Leu Cys Thr Asp Asp Tyr
 385 390 395 400
 Leu Gly Lys